



Basiliximab



Simulect®

20mg

Sterile Freeze-Dried Powder for I.V. Infusion or Injection

Immunosuppressant

DESCRIPTION AND COMPOSITION

One vial of Basiliximab (Simulect®) 20 mg contains 20 mg basiliximab.

An ampule containing 5 mL water for injection is supplied for dissolution.

Pharmaceutical form

Glass vials containing 20 mg sterile freeze-dried powder of basiliximab for intravenous infusion or injection after reconstitution with 5 mL water for injection.

Active substance

Basiliximab

Excipients

A vial of Basiliximab (Simulect®) contains, in addition to basiliximab, potassium dihydrogen phosphate, disodium phosphate, anhydrous, sodium chloride, sucrose, mannitol, and glycine. A solvent ampule contains water for injection. No preservatives are included.

INDICATIONS

Basiliximab (Simulect®) is indicated for the prophylaxis of acute organ rejection in *de novo* renal transplantation in adult and pediatric patients. It is to be used concomitantly with ciclosporin for microemulsion- and corticosteroid-based immunosuppression or in a triple maintenance immunosuppressive regimen containing ciclosporin for microemulsion, corticosteroids and either azathioprine or mycophenolate mofetil.

DOSAGE AND ADMINISTRATION

Dosage

General target population:

Adults

The standard total dose is 40 mg, given in two doses of 20 mg each. The first 20 mg dose should be given within 2 hours prior to transplantation surgery. Basiliximab (Simulect[®]) must not be administered unless it is absolutely certain that the patient will receive the graft and concomitant immunosuppression. The second 20 mg dose should be given 4 days after transplantation. The second dose should be withheld if severe hypersensitivity reactions to Basiliximab (Simulect[®]) or graft loss occur (see section WARNINGS AND PRECAUTIONS).

Special populations:

Pediatric patients (1-17 years)

In pediatric patients weighing less than 35 kg, the recommended total dose is 20 mg, given in two doses of 10 mg each. In pediatric patients weighing 35 kg or more, the recommended dose is the adult dose, i.e. a total dose of 40 mg, given in two doses of 20 mg each. The first dose should be given within 2 hours prior to transplantation surgery. Basiliximab (Simulect[®]) must not be administered unless it is absolutely certain that the patient will receive the graft and concomitant immunosuppression. The second dose should be given 4 days after transplantation. The second dose should be withheld if severe hypersensitivity reactions to Basiliximab (Simulect[®]) or graft loss occur (see section WARNINGS AND PRECAUTIONS).

Geriatric patients (≥ 65 years)

There are limited data available on the use of Basiliximab (Simulect[®]) in the elderly, but there is no evidence that elderly patients require a different dosage from younger adult patients.

Method of administration

Reconstituted Basiliximab (Simulect[®]) can be administered either as an intravenous infusion over 20-30 minutes or as a bolus injection.

For information on reconstituting Basiliximab (Simulect[®]), see section INSTRUCTIONS FOR USE AND HANDLING.

CONTRAINDICATIONS

Basiliximab (Simulect[®]) is contraindicated in patients with known hypersensitivity to basiliximab or any other component of the formulation (see section DESCRIPTION AND COMPOSITION/ Excipients).

WARNINGS AND PRECAUTIONS

General

Basiliximab (Simulect[®]) should be prescribed only by physicians who are experienced in the use of immunosuppressive therapy following organ transplantation.

Patients receiving Basiliximab (Simulect[®]) should be managed in facilities equipped and staffed with adequate laboratory and supportive medical resources including medications for the treatment of severe hypersensitivity reactions.

Hypersensitivity reactions

Severe acute (less than 24 hours) hypersensitivity reactions have been observed both on initial exposure to Basiliximab (Simulect[®]) and on reexposure to a subsequent course of therapy. These included anaphylactoid type reactions such as rash, urticaria, pruritus, sneezing, wheezing, hypotension, tachycardia, dyspnoea, bronchospasm, pulmonary oedema, cardiac failure, respiratory failure and capillary leak syndrome. If severe hypersensitivity occurs, therapy with Basiliximab (Simulect[®]) should be permanently discontinued and no further dose should be administered. Caution should be exercised when patients previously given Basiliximab (Simulect[®]) are re-exposed to a subsequent course of therapy with this medicine.

There is accumulating evidence that a subgroup of patients is at increased risk of developing hypersensitivity reactions. These are patients in whom, following the initial administration of Basiliximab (Simulect[®]), the concomitant immunosuppression was discontinued prematurely due, for example, to abandoned transplantation or early loss of the graft. Acute hypersensitivity reactions were observed on re-administration of Basiliximab (Simulect[®]) for a subsequent transplantation in some of these patients.

Neoplasms and infections

Transplant patients receiving immunosuppressive regimens involving combinations with or without Basiliximab (Simulect[®]) are at increased risk of developing lymphoproliferative disorders (LPDs) (such as lymphoma) and opportunistic infections (such as cytomegalovirus, CMV). In clinical trials, the incidence of opportunistic infections was similar in patients using immunosuppressive regimens with or without Basiliximab (Simulect[®]). In a pooled analysis of two five-year extension studies, no differences were found in the incidence of malignancies and LPDs between immunosuppressive regimens with or without Basiliximab (Simulect[®]) (see section ADVERSE DRUG REACTIONS).

Vaccination

No data are available on either the effects of live and inactive vaccination or the transmission of infection by live vaccines in patients receiving Basiliximab (Simulect[®]). Nevertheless, live vaccines are not recommended for immunosuppressed patients. Inactivated vaccines may be administered to immunosuppressed patients; however, response to the vaccine may depend on the degree of the immunosuppression.

ADVERSE DRUG REACTIONS

Summary of the safety profile

Basiliximab (Simulect[®]) has been tested in four randomized, double-blind, placebo-controlled studies in renal transplant recipients: in two studies patients were concomitantly treated with ciclosporin for microemulsion and corticosteroids (346 and 380 patients), in one study patients were concomitantly treated with ciclosporin for microemulsion, azathioprine and corticosteroids (340 patients), in one study patients were concomitantly treated with ciclosporin for microemulsion, mycophenolate mofetil and corticosteroids (123 patients).

Basiliximab (Simulect[®]) has also been compared to a polyclonal anti-T-lymphocyte immunoglobulin preparation (ATG/ALG) in one active-controlled study in renal transplant recipients; all patients were concomitantly treated with ciclosporin for microemulsion, mycophenolate mofetil and corticosteroids (135 patients). Safety data in pediatric patients have been obtained from one open-label pharmacokinetic and pharmacodynamic study in renal transplant recipients (41 patients).

Incidence of Adverse Events: Basiliximab (Simulect[®]) did not appear to add to the background of adverse events seen in organ transplantation patients as a consequence of their underlying disease and the concurrent administration of immunosuppressants and other medications. In the four placebo-controlled trials, the pattern of adverse events in 590 patients treated with the recommended dose of Basiliximab (Simulect[®]) was indistinguishable from that in 595 patients treated with placebo. Basiliximab (Simulect[®]) did not increase the incidence of serious adverse events observed when compared to placebo. The overall incidence of treatment-related adverse events among all patients in the individual studies was not significantly different between the Basiliximab (Simulect[®]) (7.1% to 40%) and the placebo (7.6% to 39%) treatment groups. In the active-controlled study, fewer Basiliximab (Simulect[®]) (11.4%) than ATG/ALG (41.5%) patients experienced treatment-related adverse events.

Adult experience: The most commonly reported (>20%) events following dual or triple therapy in both treatment groups (Basiliximab (Simulect[®]) vs. Placebo or ATG/ALG) were constipation, urinary tract infection, pain, nausea, peripheral edema, hypertension, anemia, headache, hyperkalemia, hypercholesterolemia, postoperative wound complication, weight increase, increase in blood creatinine, hypophosphatemia, diarrhea, upper respiratory tract infection.

Pediatric experience: The most commonly reported (>20%) events following dual therapy in both (<35 kg vs. ≥35 kg weight) cohorts were urinary tract infection, hypertrichosis, rhinitis, pyrexia, hypertension, upper respiratory tract infection and viral infection, sepsis and constipation.

Incidence of Malignant Neoplasms: The overall incidence of malignancies among all patients in the individual studies was similar between the Basiliximab (Simulect[®]) and the comparator treatment groups. Overall, lymphoma/lymphoproliferative disease occurred in 0.1% (1/701) of patients in the Basiliximab (Simulect[®]) group compared with 0.3% (2/595) of placebo and 0% of ATG/ALG patients.

Other malignancies were reported among 1.0% (7/701) of patients in the Basiliximab (Simulect[®]) group compared with 1.2% (7/595) of placebo and 4.6% (3/65) of ATG/ALG patients.

No differences were found in the incidence of malignancies and LPDs between Basiliximab (Simulect[®]) 7% (21/295) and placebo 7% (21/291) in a pooled analysis of two five-years extension studies.

Incidence of Infectious Episodes: The overall incidence and profile of infectious episodes among dual and triple therapy patients was similar between the Basiliximab (Simulect[®]) and the placebo treatment groups (Basiliximab (Simulect[®]) = 75.9%, Placebo or ATG/ALG = 75.6%). The incidence of serious infections was similar in the Basiliximab (Simulect[®]) and comparator groups (26.1% vs. 24.8%). The incidence of CMV-infections was similar in both groups (14.6% vs. 17.3%), following either dual or triple therapy regimen.

The incidence and causes of deaths following dual or triple therapy were similar in Basiliximab (Simulect[®]) (2.9%) and placebo or ATG/ALG groups (2.6%), with the most common cause of deaths in both treatment groups being infections (Basiliximab (Simulect[®]) = 1.3%, placebo or ATG/ALG = 1.4%). In a pooled analysis of two five-year extension studies the incidence and cause of death remained similar in both treatment groups (Basiliximab (Simulect[®]) 15%, placebo 11%), the primary cause of death being cardiac-related disorders such as cardiac failure and myocardial infarction (Basiliximab (Simulect[®]) 5%, placebo 4%).

Listing of adverse drug reactions from post-marketing spontaneous reports

The following adverse drug reactions have been identified based on post-marketing spontaneous reports and are organized by system organ classes. Because these reactions are reported voluntary from a population of uncertain size, it is not always possible to reliably estimate their frequency.

Immune system disorders

Hypersensitivity/anaphylactoid reaction such as rash, urticaria, pruritus, sneezing, wheezing, bronchospasm, dyspnea, pulmonary edema, cardiac failure, hypotension, tachycardia, respiratory failure, capillary leak syndrome, Cytokine release syndrome.

INTERACTIONS

Because Basiliximab (Simulect[®]) is an immunoglobulin, no metabolic drug-drug interactions are to be expected.

Concomitant medications routinely administered in organ transplantation

In addition to ciclosporin for microemulsion, steroids, azathioprine and mycophenolate mofetil, other concomitant medications routinely administered in organ transplantation have been administered in clinical trials without any incremental adverse reactions. These concomitant medications include systemic antiviral, antibacterial and antimycotic medications, analgesics, antihypertensive medications such as beta-blocking agents or calcium channel blockers, and diuretics.

In the original phase 3 studies during the first 3 months post-transplantation, 14% of patients in the Basiliximab (Simulect[®]) group and 27% of patients in the placebo group had an acute

rejection episode treated with antibody therapy (OKT 3 or ATG/ALG), with no increase in adverse events or infections in the Basiliximab (Simulect®) group as compared to placebo.

Three clinical trials have investigated Basiliximab (Simulect®) use in combination with a triple therapy regimen which included either azathioprine or mycophenolate mofetil. The total body clearance of Basiliximab (Simulect®) was reduced by an average 22% when azathioprine was added to a regimen consisting of ciclosporin for microemulsion and corticosteroids. The total body clearance of Basiliximab (Simulect®) was reduced by an average 51% when mycophenolate mofetil was added to a regimen consisting of ciclosporin for microemulsion and corticosteroids. The use of Basiliximab (Simulect®) in a triple therapy regimen including azathioprine or mycophenolate mofetil did not increase adverse events or infections in the Basiliximab (Simulect®) group as compared to placebo (see section ADVERSE DRUG REACTIONS).

Human antimurine antibody (HAMA)

Human antimurine antibody (HAMA) responses were reported in a clinical trial of 172 patients treated with Basiliximab (Simulect®), without predictive value for clinical tolerability. The incidence was 2/138 in patients not exposed to muromonab-CD3 and 4/34 in patients who received muromonab-CD3 concomitantly. The use of Basiliximab (Simulect®) does not preclude subsequent treatment with murine antilymphocyte antibody preparations.

WOMEN OF CHILDBEARING POTENTIAL, PREGNANCY, BREAST-FEEDING AND FERTILITY

Women of childbearing potential

Women of child-bearing potential should use adequate contraception to prevent pregnancy and continue its use for an additional 4 months after the last dose of Basiliximab (Simulect®).

Pregnancy

There is no adequate information for use in pregnant women. Basiliximab (Simulect®) should not be given to pregnant women except in cases where the potential benefit for the mother outweighs the potential risk for the fetus.

Breast-feeding

There is no animal or human data available concerning excretion of basiliximab into breast milk. However, since Basiliximab (Simulect®) is an immunoglobulin G (IgG_{1K}) antibody, it may cross the human placenta and may be excreted in human milk.

Women receiving Basiliximab (Simulect®) should not breastfeed for 4 months following the last dose.

Fertility

No human data on the effect of basiliximab on fertility are available. Formal studies of the potential effect of Basiliximab (Simulect®) on animal fertility have not been conducted (see section NON CLINICAL SAFETY DATA).

OVERDOSAGE

In clinical studies Basiliximab (Simulect[®]) has been administered to humans in single doses of up to 60 mg and multiple doses of up to 150 mg over 24 days with no untoward acute effects.

CLINICAL PHARMACOLOGY

Pharmacodynamic properties (PD)

Basiliximab (Simulect[®]) is a murine/human chimeric monoclonal antibody (IgG_{1κ}) that is directed against the interleukin-2 receptor alpha-chain (CD25 antigen), which is expressed on the surface of T-lymphocytes in response to antigenic challenge. Basiliximab (Simulect[®]) specifically binds with high affinity (K_D-value 0.1 nM) to the CD25 antigen on activated T-lymphocytes expressing the high affinity interleukin-2 receptor and thereby prevents binding of interleukin-2, the signal for T-cell proliferation. Complete and consistent blocking of the interleukin-2 receptor is maintained as long as serum basiliximab levels exceed 0.2 micrograms/mL. As concentrations fall below this level, expression of the CD25 antigen returns to pretherapy values within 1 to 2 weeks. Basiliximab (Simulect[®]) does not cause myelosuppression.

Pharmacokinetic properties (PK)

Single-dose and multiple-dose pharmacokinetic studies have been conducted in patients undergoing kidney transplantation. Cumulative doses ranged from 15 mg up to 150 mg.

Absorption

Peak serum concentration following intravenous infusion of 20 mg over 30 minutes is 7.1 ± 5.1 mg/L. There is a proportional increase in C_{max} and AUC with dose up to the highest tested single dose of 60 mg.

Distribution

The volume of distribution at steady state is 8.6 ± 4.1 L. The extent and degree of distribution to various body compartments have not been fully studied. *In vitro* studies using human tissues indicate that Basiliximab (Simulect[®]) binds only to lymphocytes and macrophages/monocytes.

Metabolism

Not applicable.

Elimination

The terminal half-life is 7.2 ± 3.2 days. Total body clearance is 41 ± 19 mL/h.

Characteristics in patients

No clinically relevant influence of body weight or gender on distribution volume or clearance has been observed in adult patients. Elimination half-life was not influenced by age (20 to 69 years), gender or race.

Disposition in adult liver transplant patients is characterized by a steady-state distribution volume of 7.5 ± 2.5 L, half-life of 4.1 ± 2.1 days and clearance of 75 ± 24 mL/h. Contributing to clearance were drug loss via drained ascites fluid and post-operative bleeding. Offsetting the faster drug clearance was a lower receptor-saturating concentration threshold of 0.1

micrograms/mL in this population. Hence, the duration of IL-2Ralpha blockade at a given Basiliximab (Simulect®) dose level is similar to that seen in adult renal transplant patients.

Pediatrics

The pharmacokinetics of Basiliximab (Simulect®) were assessed in 39 pediatric *de novo* renal transplantation patients. In infants and children (age 1 to 11 years, n=25), the steady-state distribution volume was 4.8 ± 2.1 L, half-life was 9.5 ± 4.5 days and clearance was 17 ± 6 mL/h. Distribution volume and clearance are reduced by about 50% compared to adult renal transplantation patients. Disposition parameters were not influenced to a clinically relevant extent by age (1 to 11 years), body weight (9–37 kg) or body surface area (0.44 to 1.20 m²) in this age group. In adolescents (age 12 to 16 years, n=14), the steady-state distribution volume was 7.8 ± 5.1 L, half-life was 9.1 ± 3.9 days and clearance was 31 ± 19 mL/h. Disposition in adolescents was similar to that in adult renal transplantation patients. The relationship between serum concentration and receptor saturation was assessed in 13 patients and was similar to that characterized in adult renal transplantation patients.

CLINICAL STUDIES

The efficacy of Basiliximab (Simulect®) in prophylaxis of organ rejection in *de novo* renal transplantation has been demonstrated in double-blind placebo-controlled studies. Results from two pivotal 12-month multicenter studies comparing Basiliximab (Simulect®) with placebo show that Basiliximab (Simulect®), used concomitantly with ciclosporin for microemulsion and corticosteroids, significantly reduces the incidence of acute rejection episodes both within 6 (31% vs. 45%, $p < 0.001$) and 12 (33% vs. 48%, $p < 0.001$) months after transplantation. There was no significant difference between Basiliximab (Simulect®) and placebo treated patients in graft survival after 6 and 12 months (at 12 months 32 graft losses on Basiliximab (Simulect®) (9%) and 37 graft losses on placebo (10%)). The incidence of acute rejection episode was substantially lower in patients receiving Basiliximab (Simulect®) and a triple drug immunosuppressive regimen.

Results from two multicenter double-blind studies comparing Basiliximab (Simulect®) with placebo show that Basiliximab (Simulect®) significantly reduces the incidence of acute rejection episodes within 6 months after transplantation when used concomitantly with ciclosporin for microemulsion, corticosteroids, and either azathioprine (21% vs. 35%, $p=0.005$ Fisher's exact) or mycophenolate mofetil (15% vs. 27%, $p=0.046$ K-M). Graft loss occurred in 6% of Basiliximab (Simulect®) and 10% of placebo patients by 6 months. The adverse event profile remained comparable between treatment groups.

One 12-month active-controlled randomized open-label study compared Basiliximab (Simulect®) used concomitantly with early ciclosporin for microemulsion to a polyclonal anti-T-lymphocyte immunoglobulin preparation (ATG/ALG) with delayed ciclosporin for microemulsion. Both groups received corticosteroids and mycophenolate mofetil. Biopsy proven rejection occurred in 19% of Basiliximab (Simulect®) and 20% of ATG/ALG treated patients within 12 months post-transplant.

In a pooled analysis of two five-year open-label extension studies (586 patients total) the combined graft and patient survival rates were not statistically different for the Basiliximab (Simulect®) and placebo groups. Extension studies also showed that patients who experienced an acute rejection episode during the first year after transplantation experienced more graft

losses and deaths over the five-year follow-up period than patients who had no rejection. These events were not influenced by Basiliximab (Simulect®).

Basiliximab (Simulect®) was used concomitantly with ciclosporin for microemulsion and steroids in an uncontrolled study in pediatric *de novo* renal transplant recipients. Acute rejection occurred in 14.6% of patients by 6 months post-transplantation, and in 24.3% by 12 months. Overall the adverse event profile was consistent with general clinical experience in the pediatric renal transplantation population and with the profile in the controlled adult transplantation studies.

Of 339 renal transplant patients treated with Basiliximab (Simulect®) and tested for anti-idiotypic antibodies, 4 (1.2%) developed an anti-idiotypic antibody response. In a clinical trial with 172 patients receiving Basiliximab (Simulect®), the incidence of human anti-murine antibody (HAMA) in renal transplantation patients treated with Basiliximab (Simulect®) was 2/138 in patients not exposed to muromonab-CD3 and 4/34 in patients who received muromonab-CD3 concomitantly. The available clinical data on the use of muromonab-CD3 in patients previously treated with Basiliximab (Simulect®) suggest that subsequent use of muromonab-CD3 or other murine anti-lymphocytic antibody preparations is not precluded.

NON-CLINICAL SAFETY DATA

No local irritation potential was observed in a sensitive rabbit model intravenously injected with up to 4 mg/mL of basiliximab.

No toxicity was observed when rhesus monkeys received intravenous doses of either up to 5 mg/kg basiliximab twice weekly for 4 weeks followed by an 8-week withdrawal period or 24 mg/kg basiliximab weekly for 39 weeks followed by a 13-week withdrawal period. The highest dose resulted in approximately 1000 times the systemic exposure (AUC) observed in renal transplant patients given the recommended clinical dose together with concomitant immunosuppressive therapy.

No maternal toxicity, embryotoxicity, or teratogenicity was observed in cynomolgous monkeys 100 days *post coitum* following intravenous bolus injections of up to 5 mg/kg basiliximab administered twice weekly during the organogenesis period.

Formal non-clinical studies on the potential effects of basiliximab on fertility have not been conducted (see section WOMEN OF CHILDBEARING POTENTIAL, PREGNANCY, BREAST-FEEDING AND FERTILITY).

No mutagenic potential was observed *in vitro*.

INCOMPATIBILITIES

No known incompatibilities.

STORAGE

Shipping and storage should be under refrigerated conditions (2 to 8°C). See folding box.

Basiliximab (Simulect®) should not be used after the date marked “EXP” on the pack.

Basiliximab (Simulect®) should be kept out of the reach and sight of children.

INSTRUCTIONS FOR USE AND HANDLING

Basiliximab (Simulect®) 20 mg

To prepare the infusion/injection solution, add 5 mL of water for injection from the accompanying ampule aseptically to the vial containing the Basiliximab (Simulect®) powder. Shake the vial gently to dissolve the powder. Use the reconstituted, colourless, clear to opalescent solution as soon as possible, but it may be stored at 2 to 8°C for 24 hours or at room temperature for 4 hours. Discard the reconstituted solution if not used within 24 hours.

The reconstituted solution is isotonic and may be given as a bolus injection or diluted to a volume of 50 mL or greater with normal saline or dextrose 5% for infusion.

Since no data are available on the compatibility of Basiliximab (Simulect®) with other intravenous substances, Basiliximab (Simulect®) should not be mixed with other medications/substances and should always be given through a separate infusion line.

Compatibility with the following infusion sets has been verified:

Infusion bag

- Baxter minibag NaCl 0.9%

Infusion sets

- Luer Lock™, H. Noolens
- Sterile vented i.v. set, Abbott
- Infusion set, Codan
- Infusomat™, Braun
- Infusionsgerät R 87 plus, Ohmeda
- Lifecare 5000™ Plumset Microdrip, Abbott
- Vented basic set, Baxter
- Flashball device, Baxter
- Vented primary administration set, Imed

Compatibility with other commercial devices has not been tested.

AVAILABILITY

20 ml vial and one ampule containing 5 ml Water for Injection supplied for dissolution

CAUTION

Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph

The patient is advised to seek IMMEDIATE medical attention at the first sign of adverse drug reaction.

Registration Number/Date of Renewal of Authorization:

20 mg Sterile Freeze-Dried Powder for I.V. Infusion/ Injection: BR-1051 / 09 Feb 2015

Manufactured by:**Novartis Pharma Stein AG**

Schaffhauserstrasse, 4332 Stein, Switzerland

Imported by:**Novartis Healthcare Philippines, Inc.**

5th and 6th Floors Ayala North Exchange Building, Tower 1,

Ayala Avenue corner Salcedo & Amorsolo Sts.,

Brgy. San Lorenzo, Makati, Metro Manila

Information issued: January 2014

® = registered trademark

Novartis Pharma AG, Basel, Switzerland