

ASCORBIC ACID

C-4 KIDS
100mg / 5mL Syrup
VITAMIN

FORMULATION:

Each 5 mL (1 teaspoonful) syrup contains:
Ascorbic acid (equivalent to Sodium
Ascorbate 112.5 mg), USP 100 mg

DESCRIPTION:

Clear, orange syrup with orange flavor.

WHAT IS IN THE MEDICINE?

Vitamin C, is a water-soluble vitamin, is essential for the synthesis of collagen and intercellular material.

STRENGTH OF THE MEDICINE.

Please see formulation

WHAT IS THIS MEDICINE USED FOR?

Sodium ascorbate used to increase body resistance against stress, colds and other infections. Also for the prevention and treatment of scurvy.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

Children 9-13 years. old: 5-10 mL
(1-2 teaspoonful)
4-8 years old: 5 mL (1 teaspoonful)
1-3 years old: 2.5 mL (½ teaspoonful)

To be taken once daily for supplementation, twice or thrice a daily for therapeutic use. Or as prescribed by the physician.

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

If you are allergic to any ingredients.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE?

Large doses may result in hyperoxaluria and the formation of renal calcium oxalate calculi and sodium ascorbate should therefore be given with care to patients with hyperoxaluria.

This product contains FD & C #5 (Tartrazine) which may cause allergic reactions including bronchial asthma in certain susceptible persons.

UNDESIRABLE EFFECTS OF THIS MEDICINE?

Large doses of sodium ascorbate have resulted in haemolysis in patients with G6PD deficiency.

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Sodium ascorbate may increase the absorption of iron in iron-deficiency states.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, take it as soon as you remember.

SIGNS AND SYMPTOMS OF OVERDOSE

Administration of high doses may cause restlessness, insomnia, headache, esophagitis and diarrhea.

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

Seek emergency medical attention.

HOW SHOULD YOU KEEP THIS MEDICINE?

Store at temperatures not exceeding 30°C.

Protect from light.

Always keep containers tightly closed.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

If symptoms persist consult your doctor immediately.

AVAILABILITY:

Boston Round Amber Bottle x 60 mL and 120 mL (Box of 1's)

Manufactured by:
Drugmaker's Laboratories, Inc.
E & E Industrial Complex,
Brgy. San Antonio, San Pedro, Laguna
For:



Terramedic
INCORPORATED
E & E Industrial Complex, Narra Rd.,
Brgy. San Antonio, San Pedro, Laguna

For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph

Registration No.: DR-XY44350

Date of First Authorization / Renewal of Authorization:
March 19, 2015/ April 30, 2018

The Patient Information Leaflet (PIL) was last revised in 06/2022