


확	확인자	QC (문안)
인	박은영	안재윤
날짜	2020. 02. 07	2020. 02. 07

 <b>DW PKG 인쇄사양서</b>	
product	
제품명	디옥타헨탁액 설명서 필리핀수출용(일반의약품)
포장단위	
size (가로x세로)	
115 X 220 mm	
종이재질	
모조-80g/m <sup>2</sup>	
코팅 (CR, 라미레이팅, UV 등)	
후가공	
접착 (맞뚜껑, 1면, 1면자동 등)	
특이사항	
문안 변경	
colors	
	BLACK
자재코드	
2000316	
에디션 넘버	
H01	
바코드 (13자리)	
제조사	
(주)대웅제약/경기도 화성시 향남읍 제약공단4길 35-14	
판매원	

## DIOCTAHEDRAL SMECTITE

### DEHECTA

150mcg/mL (3g/20mL) Oral suspension  
*Antidiarrheal*

#### Formulation

Each sachet (20ml) contains: Dioctahedral Smectite ..... 3g

#### Description

Light to dark pink suspension with vanilla and strawberry flavor.

#### Indications

- Symptomatic treatment of painful symptoms associated with oesophago-gastrointestinal diseases in adults.
- Acute and chronic diarrhea in adults.
- Acute diarrhea in children over 24 months.

#### Dosage and Administration

##### Adults:

Take 20ml 3 times a day.

For the first 3 days of acute diarrhea, daily dose can be doubled.

For esophagitis, take after meals; for other indications between meals.

##### Children (over 24 months):

Take 40-60ml in 3 divided doses per day for 3 days.

Followed by 40ml in 3 divided doses per day for 4 days.

Shake well before use.

#### Contraindications

- Children under 24 months.
- Pregnant women, women who are likely to be pregnant, and lactating women.

#### Precautions

- Patients taking other drugs: This drug has adsorbent property and if administered with other drugs it may affect absorption or absorption time that be taken at intervals.
- Patients with severe chronic constipation.
- Patients with a history of intestinal obstruction.

#### Warnings

- Follow the prescribed usage and dosage.
- When this drug is taken by children, take under the guidance and supervision of a guardian.
- This drug is not a rehydration therapy that in case of dehydration, provide adequate fluid supply.
- If symptoms did not improve after 7 days of taking, immediately stop taking this medicine and consult with your doctor or pharmacist.

#### Interactions

As the adsorbent property of this product may interfere with the rates and/or levels of absorption of other agents, it is recommended to treat with other drugs in time intervals.

#### Adverse Events

- Digestive system: bloating, vomiting, constipation, etc.
- Skin: allergic symptoms such as urticaria, rash, itching and angioedema.

#### Storage Conditions

- Store at temperature not exceeding 30°C.
- Keep out of reach of children.
- Storing out of their original containers may cause accidents or deterioration of quality of medicines. Place in the original container and keep tightly closed.

#### Caution

Foods, Drugs, Devices, and Cosmetics Act prohibit dispensing without prescription.

For suspected adverse drug reaction, report to the FDA: [www.fda.gov.ph](http://www.fda.gov.ph).

Seek medical attention immediately at the first sign of any adverse drug reaction.

#### Availability

20ml in PET/PE/AL/SUR/LLDPE Sachet (Box of 20's).

#### Shelf-life

36 months

FDA Registration No.: DR-XY44417

Date of First Authorization: 08 April 2015

Date of Last Revision: February 2020

Manufactured by:

 **DAEWOONG**  
 PHARMACEUTICAL CO., LTD.

35-14, Jeyakongdan 4-gil, Hyangnam-eup,  
 Hwaseong-si, Gyeonggi-do, Korea

Imported & Distributed by:

 **DAEWOONG**  
 PHARMA PHILIPPINES INC.

Unit 1701, 17th Floor, One World Place Condominium,  
 32nd Street, Bonifacio Global City Fort Bonifacio  
 Taguig, Metro Manila

H01  
 2000316