

Product	Ferinject Singlepack Vials	
Country	Philippines	
Manufacturer	VIT	
SAP Number	3004585-04	Min. Ver.: 04
Manufacturer Identification Number	n/a	
Fonts	Arial (min. 9pt)	
Dimensions	148x900 mm	

Colours/Flats
Black

Non-Printed Colours
Dye Cut

RC 10/11/2022
Approved by KRC

Ferric Carboxymaltose

ferinject®



**50 mg/mL Solution for Injection/Infusion (I.V.)
Anti-anemia**

1. NAME OF THE MEDICINAL PRODUCT

Ferric carboxymaltose (Ferinject) 50 mg/mL Solution for injection/infusion.

2. FORMULATION/COMPOSITION

Iron (as ferric carboxymaltose), Ph.Eur, USP..... 50mg

Excipient(s) with known effect

One mL of solution contains up to 5.5 mg (0.24 mmol) sodium, see section 4.4. For the full list of excipients, see section 6.1.

3. PRODUCT DESCRIPTION

Solution for injection/infusion. Dark brown, non-transparent, aqueous solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Ferric carboxymaltose (Ferinject) is indicated for the treatment of iron deficiency when (see section 5.1):

- oral iron preparations are ineffective.
- oral iron preparations cannot be used.
- there is a clinical need to deliver iron rapidly.

The diagnosis of iron deficiency must be based on laboratory tests.

4.2 Posology and method of administration

Monitor carefully patients for signs and symptoms of hypersensitivity reactions during and following each administration of ferric carboxymaltose (Ferinject).

Ferric carboxymaltose (Ferinject) should only be administered when staff trained to evaluate and manage anaphylactic reactions is immediately available, in an environment where full resuscitation facilities can be assured. The patient should be observed for adverse effects for at least 30 minutes following each ferric carboxymaltose (Ferinject) administration (see section 4.4).

Posology

The posology of ferric carboxymaltose (Ferinject) follows a stepwise approach: [1] determination of the individual iron need, [2] calculation and administration of the iron dose(s), and [3] post-iron repletion assessments. These steps are outlined below:

Step 1: Determination of the iron need

The individual iron need for repletion using ferric carboxymaltose (Ferinject) is determined based on the patient's body weight and haemoglobin (Hb) level. Refer to Table 1 for determination of the iron need:

Table 1: Determination of the iron need

g/dL	mmol/L	Patient body weight		
		below 35 kg	35 kg to <70 kg	70 kg and above
<10	<6.2	500 mg	1,500 mg	2,000 mg
10 to <14	6.2 to <8.7	500 mg	1,000 mg	1,500 mg
≥14	≥8.7	500 mg	500 mg	500 mg

Iron deficiency must be confirmed by laboratory tests as stated in 4.1.

Step 2: Calculation and administration of the maximum individual iron dose(s)

Based on the iron need determined above the appropriate dose(s) of Ferric carboxymaltose (Ferinject) should be administered taking into consideration the following:

- A single Ferric carboxymaltose (Ferinject) administration should not exceed:
 - 15 mg iron/kg body weight (for administration by intravenous injection) or 20 mg iron/kg body weight (for administration by intravenous infusion)
 - 1,000 mg of iron (20 mL Ferric carboxymaltose (Ferinject))

The maximum recommended cumulative dose of Ferric carboxymaltose (Ferinject) is 1,000 mg of iron (20 mL Ferric carboxymaltose (Ferinject)) per week.

Step 3: Post-iron repletion assessments

Re-assessment should be performed by the clinician based on the individual patient's condition. The Hb level should be re-assessed no earlier than 4 weeks post final Ferrinjection administration to allow adequate time for erythropoiesis and iron utilisation. In the event the patient requires further iron repletion, the iron need should be recalculated using Table 1 above. (See section 5.1.)

Special Population – patients with haemodialysis-dependent chronic kidney disease

A single maximum daily dose of 200 mg iron should not be exceeded in haemodialysis-dependent chronic kidney disease patients (see also section 4.4).

Paediatric population

The use of Ferric carboxymaltose (Ferinject) has not been studied in children, and therefore is not recommended in children under 14 years.

Method of administration

Ferric carboxymaltose (Ferinject) must only be administered by the intravenous route:

- by injection, or
- by infusion, or
- during a haemodialysis session undiluted directly into the venous limb of the dialyser.

Ferric carboxymaltose (Ferinject) must not be administered by the subcutaneous or intramuscular route.

Intravenous injection

Ferric carboxymaltose (Ferinject) may be administered by intravenous injection using undiluted solution. The maximum single dose is 15 mg iron/kg body weight but should not exceed 1,000 mg iron. The administration rates are as shown in Table 2:

Table 2: Administration rates for intravenous injection of Ferric carboxymaltose (Ferinject)

Volume of Ferric carboxymaltose (Ferinject) required	Equivalent iron dose	Administration rate / Minimum administration time
2 to 4 mL	100 to 200 mg	No minimal prescribed time
>4 to 10 mL	>200 to 500 mg	100 mg iron / min
>10 to 20 mL	>500 to 1,000 mg	15 minutes

Intravenous infusion

Ferric carboxymaltose (Ferinject) may be administered by intravenous infusion, in which case it must be diluted. The maximum single dose is 20 mg iron/kg body weight, but should not exceed 1,000 mg iron.

For infusion, Ferric carboxymaltose (Ferinject) must only be diluted in sterile 0.9% m/V sodium chloride solution as shown in Table 3. Note: for stability reasons, Ferric carboxymaltose (Ferinject) should not be diluted to concentrations less than 2 mg iron/mL (not including the volume of the ferric carboxymaltose solution). For further instructions on dilution of the medicinal product before administration, see section 6.6.

Table 3: Dilution plan of Ferric carboxymaltose (Ferinject) for intravenous infusion

Volume of Ferric carboxymaltose (Ferinject) required	Equivalent iron dose	Maximum amount of sterile 0.9% m/V sodium chloride solution	Minimum administration time
2 to 4 mL	100 to 200 mg	50 mL	No minimal prescribed time
>4 to 10 mL	>200 to 500 mg	100 mL	6 minutes
>10 to 20 mL	>500 to 1,000 mg	250 mL	15 minutes

4.3 Contraindications

The use of Ferric carboxymaltose (Ferinject) is contraindicated in cases of:

- hypersensitivity to the active substance, to Ferric carboxymaltose (Ferinject) or any of its excipients listed in section 6.1.
- known serious hypersensitivity to other parenteral iron products.
- anaemia not attributed to iron deficiency, e.g. other microcytic anaemia.
- evidence of iron overload or disturbances in the utilisation of iron.

4.4 Special warnings and precautions for use

Hypersensitivity reactions

Parenterally administered iron preparations can cause hypersensitivity reactions including serious and potentially fatal anaphylactic/anaphylactoid reactions. Hypersensitivity reactions have also been reported after previously uneventful doses of parenteral iron complexes. There have been reports of hypersensitivity reactions which progressed to Kounis syndrome (acute allergic coronary arteriospasm that can result in myocardial infarction, see section 4.8).

The risk is enhanced for patients with known allergies including drug allergies, including patients with a history of severe asthma, eczema or other atopic allergy.

There is also an increased risk of hypersensitivity reactions to parenteral iron complexes in patients with immune or inflammatory conditions (e.g. systemic lupus erythematosus, rheumatoid arthritis).

Ferric carboxymaltose (Ferinject) should only be administered when staff trained to evaluate and manage anaphylactic reactions are immediately available, in an environment where full resuscitation facilities can be assured. Each patient should be observed for adverse effects for at least 30 minutes following each Ferric carboxymaltose (Ferinject) administration. If hypersensitivity reactions or signs of intolerance occur during administration, the treatment must be stopped immediately. Facilities for cardio respiratory resuscitation and equipment for handling acute anaphylactic/anaphylactoid reactions should be available, including an injectable 1:1000 adrenaline solution. Additional treatment with antihistamines and/or corticosteroids should be given as appropriate.

Hypophosphataemia / Hypophosphataemic osteomalacia

Parenterally iron preparations can lead to hypophosphataemia which is transient and without clinical symptoms. Hypophosphataemia requiring treatment has mainly been reported in individual cases, in patients with known risk factors and after sustained higher dosing.

Cases of symptomatic hypophosphataemia leading to hypophosphataemic osteomalacia, and fractures requiring clinical intervention, including surgery, were reported after market introduction. In case of arthralgia or bone pain, patients should be advised to seek medical advice.

Patients receiving multiple higher doses as part of long-term treatment, and who have underlying risk factors (e.g. vitamin D deficiency, calcium and phosphate malabsorption, secondary hyperparathyroidism, hereditary haemorrhagic telangiectasia, inflammatory bowel disease and osteoporosis) should be monitored for hypophosphataemic osteomalacia, including serum phosphate control. In case of persistent hypophosphataemia, treatment with Ferrinjection should be re-evaluated.

Hepatic or renal impairment

In patients with liver dysfunction, parenteral iron should only be administered after careful benefit/risk assessment. Parenteral iron administration should be avoided in patients with hepatic dysfunction where iron overload is a precipitating factor, in particular Porphyria Cutanea Tarda (PCT). Careful monitoring of iron status is recommended to avoid iron overload.

No safety data on haemodialysis-dependent chronic kidney disease patients receiving single doses of more than 200 mg iron are available.

Infection

Parenteral iron must be used with caution in case of acute or chronic infection, asthma, eczema or atopic allergies. It is recommended that the treatment with Ferric carboxymaltose (Ferinject) is stopped in patients with ongoing bacteraemia. Therefore, in patients with chronic infection a benefit/risk evaluation has to be performed, taking into account the suppression of erythropoiesis.

Extravasation

Caution should be exercised to avoid paravenous leakage when administering Ferric carboxymaltose (Ferinject). Paravenous leakage of Ferric carboxymaltose (Ferinject) at the administration site may lead to irritation of the skin and potentially long lasting brown discoloration at the site of administration. In case of paravenous leakage, the administration of Ferric carboxymaltose (Ferinject) must be stopped immediately.

Excipients

One mL of undiluted Ferric carboxymaltose (Ferinject) contains up to 5.5 mg (0.24 mmol) of sodium. This has to be taken into account in patients on a sodium-controlled diet.

Paediatric population

The use of Ferric carboxymaltose (Ferinject) has not been studied in children.

4.5 Interaction with other medicinal products and other forms of interaction

The absorption of oral iron is reduced when administered concomitantly with parenteral iron preparations. Therefore, if required, oral iron therapy should not be started for at least 5 days after the last administration of Ferric carboxymaltose (Ferinject).

4.6 Fertility, pregnancy and lactation

Pregnancy

There are limited data from the use of Ferric carboxymaltose (Ferinject) in pregnant women (see section 5.1). A careful benefit/risk evaluation is required before use during pregnancy and Ferric carboxymaltose (Ferinject) should not be used during pregnancy unless clearly necessary.

Iron deficiency occurring in the first trimester of pregnancy can in many cases be treated with oral iron. Treatment with Ferric carboxymaltose (Ferinject) should be confined to the second and third trimester if the benefit is judged to outweigh the potential risk for both the mother and the foetus.

Foetal bradycardia may occur following administration of parenteral irons. It is usually transient and a consequence of a hypersensitivity reaction in the mother. The unborn baby should be carefully monitored during intravenous administration of parenteral irons to pregnant women.

Animal data suggest that iron released from Ferric carboxymaltose (Ferinject) can cross the placental barrier and that its use during pregnancy may influence skeletal development in the fetus (see section 5.3).

Breast-feeding

Clinical studies showed that transfer of iron from Ferric carboxymaltose (Ferinject) to human milk was negligible (<1%). Based on limited data on breast-feeding women it is unlikely that Ferric carboxymaltose (Ferinject) represents a risk to the breast-fed child.

Fertility

There are no data on the effect of Ferric carboxymaltose (Ferinject) on human fertility. Fertility was unaffected following Ferric carboxymaltose (Ferinject) treatment in animal studies (see section 5.3).

4.7 Effects on ability to drive and use machines

Ferric carboxymaltose (Ferinject) is unlikely to impair the ability to drive and use machines.

4.8 Adverse Drug Reactions

Table 4 presents the adverse drug reactions (ADRs) reported during clinical studies in which >8,000 subjects received Ferric carboxymaltose (Ferinject), as well as those reported from the post-marketing experience (see table footnotes for details).

The most commonly reported ADR is nausea (occurring in 2.9% of the subjects), followed by injection/infusion site reactions, hypophosphataemia, headache, flushing, dizziness and hypertension. Injection/infusion site reactions comprise several ADRs which individually are either uncommon or rare.

For subjects in clinical trials that showed a decrease in serum phosphorus, the minimum values were obtained after approximately 2 weeks, and in most cases returned to baseline values by 12 weeks following Ferric carboxymaltose (Ferinject) treatment. The most serious ADR is anaphylactoid/anaphylactic reactions (rare); fatalities have been reported. See section 4.4 for further details.

Table 4: Adverse drug reactions observed during clinical trials and post-marketing experience

System Organ Class	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Frequency not known ⁽¹⁾
Immune system disorders		Hypersensitivity	Anaphylactoid/anaphylactic reactions	
Metabolism and nutritional disorders	Hypophosphataemia			
Nervous system disorders	Headache, dizziness	Paraesthesia, dysgeusia		Loss of consciousness ⁽¹⁾
Psychiatric disorders			Anxiety ⁽²⁾	
Cardiac disorders		Tachycardia		Kounis syndrome ⁽¹⁾
Vascular disorders	Flushing, hypertension	Hypotension	Phlebitis, syncope ⁽²⁾ , presyncope ⁽²⁾	
Respiratory, thoracic and mediastinal disorders		Dyspnoea	Bronchospasm ⁽²⁾	
Gastrointestinal disorders	Nausea	Vomiting, dyspepsia, abdominal pain, constipation, diarrhoea	Flatulence	

3004585-04

System Organ Class	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Frequency not known ⁽¹⁾
Skin and subcutaneous tissue disorders		Pruritus, urticaria, erythema, rash ⁽³⁾	Angioedema ⁽²⁾ , pallor ⁽²⁾ , distant skin discoloration	Face oedema ⁽¹⁾ , dermatitis
Musculoskeletal and connective tissue disorders		Myalgia, back pain, arthralgia, pain in extremity, muscle spasms		Hypophosphataemic, osteomalacia ⁽¹⁾
General disorders and administration site conditions	Injection/infusion site reactions ⁽⁴⁾	Pyrexia, fatigue, chest pain, oedema peripheral, chills	Malaise, influenza like illness (whose onset may vary from a few hours to several days) ⁽²⁾	
Investigations		Alanine aminotransferase increased, aspartate aminotransferase increased, gamma-glutamyltransferase increased, blood lactate dehydrogenase increased, blood alkaline phosphatase increased		

- 1 ADRs exclusively reported in the post-marketing setting; estimated as rare.
2 ADRs reported in the post-marketing setting which are also observed in the clinical setting.
3 Includes the following preferred terms: rash (individual ADR determined to be uncommon) and rash erythematous, -generalised, -macular, - maculo-papular, -pruritic (all individual ADRs determined to be rare).
4 Includes, but is not limited to the following preferred terms: injection/infusion site -pain, -haematoma, -discolouration, -extravasation, -irritation, -reaction, (all individual ADRs determined to be uncommon) and -paraesthesia (individual ADR determined to be rare).
Note: ADR = Adverse drug reaction.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via:

Philippines

ADR Reporting Website: www.fda.gov.ph

4.9 Overdose and Treatment

Administration of Ferric carboxymaltose (Ferinject) in quantities exceeding the amount needed to correct iron deficit at the time of administration may lead to accumulation of iron in storage sites eventually leading to haemosiderosis. Monitoring of iron parameters such as serum ferritin and transferrin saturation may assist in recognising iron accumulation. If iron accumulation has occurred, treat according to standard medical practice, e.g. consider the use of an iron chelator.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Iron trivalent, parenteral preparation, ATC code: B03AC

Ferric carboxymaltose (Ferinject) solution for injection/infusion is a colloidal solution of the iron complex ferric carboxymaltose.

The complex is designed to provide, in a controlled way, utilisable iron for the iron transport and storage proteins in the body (transferrin and ferritin, respectively).

Red cell utilisation of ⁵⁹Fe from radio-labelled Ferric carboxymaltose (Ferinject) ranged from 91% to 99% in subjects with iron deficiency (ID) and 61% to 84% in subjects with renal anaemia at 24 days post-dose.

Ferric carboxymaltose (Ferinject) treatment results in an increase in reticulocyte count, serum ferritin levels and TSAT levels to within normal ranges.

Clinical efficacy and safety

The efficacy and safety of Ferric carboxymaltose (Ferinject) has been studied in different therapeutic areas necessitating intravenous iron to correct iron deficiency. The main studies are described in more detail below.

Cardiology

Chronic heart failure

Study CONFIRM-HF was a double-blind, randomised, 2-arm study comparing Ferric carboxymaltose (Ferinject) (n=150) vs. placebo (n=151) in subjects with chronic heart failure and ID for a treatment period of 52 weeks. At Day 1 and Week 6 (correction phase), subjects received either Ferric carboxymaltose (Ferinject) according to a simplified dosing grid using baseline Hb and body weight at screening (see section 4.2), placebo or no dose. At Weeks 12, 24, and 36 (maintenance phase) subjects received Ferric carboxymaltose (Ferinject) (500 mg iron) or placebo if serum ferritin was <100 ng/mL or 100-300 ng/mL with TSAT <20%. The treatment benefit of Ferric carboxymaltose (Ferinject) vs. placebo was demonstrated with the primary efficacy endpoint, the change in the 6- minute walk test (6MWT) from baseline to Week 24 (33±11 metres, p=0.002). This effect was sustained throughout the study to Week 52 (36±11 metres p<0.001).

Study EFFECT-HF was an open-label (with blinded endpoint evaluation), randomised, 2-arm study comparing Ferrinject (n=86) vs. standard of care (n=86) in subjects with chronic heart failure and ID for a treatment period of 24 weeks. At Day 1 and Week 6 (correction phase), subjects received either Ferrinject according to a simplified dosing grid using baseline Hb and body weight at screening (see section 4.2) or standard of care. At Week 12, (maintenance phase) subjects received Ferrinject (500 mg iron) or standard of care if serum ferritin <100 ng/ml or 100 to 300 ng/ml and TSAT <20%. The treatment benefit of Ferrinject vs. standard of care was demonstrated with the primary efficacy endpoint, the change in weight-adjusted peak VO₂ from baseline to Week 24 (LS Mean 1.04 ±0.44, p=0.02).

Nephrology

Haemodialysis-dependent chronic kidney disease

Study VIT-IV-CL-015 was an open-label, randomised parallel group study comparing Ferric carboxymaltose (Ferinject) (n=97) to iron sucrose (n=86) in subjects with ID anaemia undergoing haemodialysis. Subjects received Ferric carboxymaltose (Ferinject) or iron sucrose 2-3 times per week in single doses of 200 mg iron directly into the dialyser until the individually calculated cumulative iron dose was reached (mean cumulative dose of iron as Ferric carboxymaltose (Ferinject): 1,700 mg). The primary efficacy endpoint was the percentage of subjects reaching an increase in Hb of ≥1.0 g/dL at 4 weeks after baseline. At 4 weeks after baseline, 44.1% responded to treatment with Ferric carboxymaltose (Ferinject) (i.e. Hb increase of ≥1.0 g/dL) compared to 35.3% for iron sucrose (p=0.2254).

Non-dialysis-dependent chronic kidney disease

Study 1VIT04004 was an open-label, randomised active-control study, evaluating the safety and efficacy of Ferric carboxymaltose (Ferinject) (n=147) vs. oral iron (n=103). Subjects in the Ferric carboxymaltose (Ferinject) group received 1,000 mg of iron at baseline and 500 mg of iron at days 14 and 28, if TSAT was <30% and serum ferritin was <500 ng/mL at the respective visit. Subjects in the oral iron arm received 65 mg iron TID as ferrous sulphate from baseline to day 56. Subjects were followed-up until day 56. The primary efficacy endpoint was the percentage of subjects achieving an increase in Hb of ≥1.0 g/dL anytime between baseline and end of study or time of intervention. This was achieved by 60.54% of subjects receiving Ferric carboxymaltose (Ferinject) vs. 34.7% of subjects in the oral iron group (p<0.001). Mean haemoglobin change to day 56/end of study was 1.0 g/dL in the Ferric carboxymaltose (Ferinject) group and 0.7 g/dL in the oral iron group (p=0.034, 95% CI: 0.0, 0.7).

Gastroenterology

Inflammatory bowel disease

Study VIT-IV-CL-008 was a randomised, open-label study which compared the efficacy of Ferric carboxymaltose (Ferinject) vs. oral ferrous sulphate in reducing ID anaemia in subjects with inflammatory bowel disease (IBD). Subjects received either Ferric carboxymaltose (Ferinject) (n=111) in single doses of up to 1,000 mg iron once per week until the individually calculated iron dose (per Ganzoni formula) was reached (mean cumulative iron dose: 1,490 mg), or 100 mg iron BID as ferrous sulphate (n=49) for 12 weeks. Subjects receiving Ferric carboxymaltose (Ferinject) showed a mean increase in Hb from baseline to Week 12 of 3.83 g/dL, which was non-inferior to 12 weeks of twice daily therapy with ferrous sulphate (3.75 g/dL, p=0.8016).

Study FER-IBD-07-COR was a randomised, open-label study comparing the efficacy of Ferric carboxymaltose (Ferinject) vs. iron sucrose in subjects with remitting or mild IBD. Subjects receiving Ferric carboxymaltose (Ferinject) were dosed according to a simplified dosing grid using baseline Hb and body weight (see section 4.2) in single doses up to 1,000 mg iron, whereas subjects receiving iron sucrose were dosed according to individually calculated iron doses using the Ganzoni formula in doses of 200 mg iron until the cumulative iron dose was reached. Subjects were followed-up for 12 weeks.

65.8% of subjects receiving Ferric carboxymaltose (Ferinject) (n=240; mean cumulative iron dose: 1,414 mg) vs. 53.6% receiving iron sucrose (n=235; mean cumulative dose 1,207 mg; p=0.004) had responded at Week 12 (defined as Hb increase ≥2 g/dL). 83.8% of Ferric carboxymaltose (Ferinject)- treated subjects vs. 75.9% of iron sucrose-treated subjects achieved a Hb increase ≥2 g/dL or had Hb within normal limits at Week 12 (p=0.019).

Women's health

Post partum

Study VIT-IV-CL-009 was a randomised open-label non-inferiority study comparing the efficacy of Ferric carboxymaltose (Ferinject) (n=227) vs. ferrous sulphate (n=117) in women suffering from post-partum anaemia. Subjects received either Ferric carboxymaltose (Ferinject) in single doses of up to 1,000 mg iron until their individually calculated cumulative iron dose (per Ganzoni formula) was reached, or 100 mg of iron as oral ferrous sulphate BID for 12 weeks. Subjects were followed-up for 12 weeks. The mean change in Hb from baseline to Week 12 was 3.37 g/dL in the Ferric carboxymaltose (Ferinject) group (n=179; mean cumulative iron dose: 1,347 mg) vs. 3.29 g/dL in the ferrous sulphate group (n=89), showing non-inferiority between the treatments.

Pregnancy

Intravenous iron medicines should not be used during pregnancy unless clearly necessary. Treatment with Ferric carboxymaltose (Ferinject) should be confined to the second and third trimester if the benefit is judged to outweigh the potential risk for both the mother and the foetus, see section 4.6.

Limited safety data in pregnant women are available from study FER-ASAP-2009-01, a randomised, open-label study comparing Ferric carboxymaltose (Ferinject) (n=121) vs. oral ferrous sulphate (n=115) in pregnant women in the second and third trimester with ID anaemia for a treatment period of 12 weeks. Subjects received Ferric carboxymaltose (Ferinject) in cumulative doses of 1,000 mg or 1,500 mg of iron (mean cumulative dose: 1,029 mg iron) based on Hb and body weight at screening, or 100 mg of oral iron BID for 12 weeks. The incidence of treatment-related adverse events was similar between Ferric carboxymaltose (Ferinject) treated women and those treated with oral iron (11.4% Ferric carboxymaltose (Ferinject) group; 15.3% oral iron group). The most commonly reported treatment-related adverse events were nausea, upper abdominal pain and headache. Newborn Apgar scores as well as newborn iron parameters were similar between treatment groups.

Ferritin monitoring after replacement therapy

There is limited data from study VIT-IV-CL-008 which demonstrates that ferritin levels decrease rapidly 2-4 weeks following replacement and more slowly thereafter. The mean ferritin levels did not drop to levels where retreatment might be considered during the 12 weeks of study follow up. Thus, the available data does not clearly indicate an optimal time for ferritin retesting although assessing ferritin levels earlier than 4 weeks after replacement therapy appears premature. Thus, it is recommended that further re-assessment of ferritin should be made by the clinician based on the individual patient's condition.

5.2 Pharmacokinetic properties

Distribution

Positron emission tomography demonstrated that ⁵⁹Fe and ⁵²Fe from Ferric carboxymaltose (Ferinject) was rapidly eliminated from the blood, transferred to the bone marrow, and deposited in the liver and spleen. After administration of a single dose of Ferric carboxymaltose (Ferinject) of 100 to 1,000 mg of iron in ID subjects, maximum total serum iron levels of 37 µg/mL up to 333 µg/mL are obtained after 15 minutes to 1.21 hours respectively. The volume of the central compartment corresponds well to the volume of the plasma (approximately 3 litres).

Elimination

The iron injected or infused was rapidly cleared from the plasma, the terminal half-life ranged from 7 to 12 hours, the mean residence time (MRT) from 11 to 18 hours. Renal elimination of iron was negligible.

5.3 Preclinical safety data

Preclinical data revealed no special hazard for humans based on conventional studies of safety pharmacology, repeat dose toxicity and genotoxicity. Preclinical studies indicated that iron released from Ferric carboxymaltose (Ferinject) does cross the placental barrier and is excreted in milk in limited, controlled amounts. In reproductive toxicology studies using iron replete rabbits Ferric carboxymaltose (Ferinject) was associated with minor skeletal abnormalities in the fetus. In a fertility study in rats, there were no effects on fertility for either male or female animals. No long-term studies in animals have been performed to evaluate the carcinogenic potential of Ferric carboxymaltose (Ferinject). No evidence of allergic or immunotoxic potential has been observed. A controlled in-vivo test demonstrated no cross-reactivity of Ferric carboxymaltose (Ferinject) with anti-dextran antibodies. No local irritation or intolerance was observed after intravenous administration.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

- Sodium hydroxide (for pH adjustment)
- Hydrochloric acid (for pH adjustment)
- Water for injections

6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6. The compatibility with containers other than polyethylene and glass is not known.

6.3 Shelf life

Shelf life of the product as packaged for sale:
3 years.

Shelf life after first opening of the container:
From a microbiological point of view, preparations for parenteral administration should be used immediately.

Shelf life after dilution with sterile 0.9% m/V sodium chloride solution:

From a microbiological point of view, preparations for parenteral administration should be used immediately after dilution with sterile 0.9% m/V sodium chloride solution.

6.4 Storage Conditions

Store in the original package in order to protect from light. Do not store above 30 °C. Do not freeze. For storage conditions after dilution or first opening of the medicinal product, see section 6.3.

6.5 Dosage Form and Packaging Available

Ferric carboxymaltose (Ferinject) is supplied in a vial (type I glass) with a stopper (bromobutyl rubber) and an aluminium cap as:

- 2 mL solution containing 100 mg iron. Available in pack sizes of 1 and 5 vials.
- 10 mL solution containing 500 mg iron. Available in pack sizes of 1, 2 and 5 vials.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Inspect vials visually for sediment and damage before use. Use only those containing sediment-free, homogeneous solution.

Each vial of Ferric carboxymaltose (Ferinject) is intended for single use only. Any unused product or waste material should be disposed of in accordance with local requirements.

Ferric carboxymaltose (Ferinject) must only be mixed with sterile 0.9% m/V sodium chloride solution. No other intravenous dilution solutions and therapeutic agents should be used, as there is the potential for precipitation and/or interaction. For dilution instructions, see section 4.2.

7. MARKETING AUTHORISATION HOLDER

Imported and Distributed by:

Zuellig Pharma Corporation
Km. 14 West Service Rd. South Super Highway corner Edison Ave., Sun Valley, Parañaque City, Philippines

8. MANUFACTURER

Manufactured by:

IDT Biologika GmbH
Am Pharmapark
D-06861 Dessau-Rosslau
Germany

Manufactured for:

Vifor (International) Inc.
Rechenstrasse 37
9014 St.Gallen
Switzerland

9. CAUTION STATEMENT

Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

10. ADR REPORTING STATEMENT

For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph. Seek medical attention immediately at the first sign of any adverse drug reaction.

11. REGISTRATION NUMBER

DR-XY46099

12. DATE OF FIRST AUTHORISATION

13 OCT 2017

13. DATE OF REVISION OF THE TEXT

02 MAR 2022

3004585-04