

# ORAL REHYDRATION SALTS

## SODALITE®-75 Powder for Oral Solution REHYDRATION SOLUTIONS

### FORMULATION:

Each sachet contains	
Glucose, Anhydrous USP.....	2.70 g
Trisodium Citrate Dihydrate USP.....	580 mg
Sodium Chloride USP.....	520 mg
Potassium Chloride USP.....	300 mg

### Equivalent in mmol/L

Glucose.....	75	Potassium.....	20
Sodium.....	75	Citrate.....	10
Chloride.....	65	(or 30 mmol bicarbonate)	

### PRODUCT DESCRIPTION:

SODALITE®-75 is an ORS which appears as white crystalline powder in a sachet, when reconstituted it is colorless liquid in orange flavor.

### WHAT IS IN THE MEDICINE?

This medicine conforms with the WHO reduced osmolarity glucose-based oral rehydration salts (ORS), the preferred method for treating dehydration.

### STRENGTH OF THE MEDICINE

See formulation above

### WHAT IS THE MEDICINE USED FOR?

This product is used for the treatment of dehydration due to diarrhea.

### HOW MUCH AND HOW OFTEN SHOULD YOU USE THE MEDICINE?

For Children: Less than 2 years old -- ½ glass (50 to 100 mL)  
Adult: 1 to 2 glasses. Or as prescribed by the physician.

### DIRECTION FOR USE:

Dissolve total contents completely in 200 mL of cooled boiled water.

### WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

If you are allergic to any ingredient of the product or if you have glucose malabsorption.

### UNDESIRABLE EFFECTS

There are no expected undesirable effects if the recommended dilution, volume of ORS solution to be administered and dosage are followed.

### WHAT OTHER MEDICINE OR FOODS SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

This medicine is not expected to interact with food or medicines.

### WHAT SHOULD YOU DO IF YOU MISS A DOSE?

Just take the next dose if still needed to help replace fluids and electrolytes lost due to diarrhea and/or vomiting.

### HOW SHOULD YOU KEEP THE MEDICINE?

- Keep the product out of reach and sight of children.
- Store at temperatures not exceeding 30°C

### SIGNS AND SYMPTOMS OF OVERDOSE

There is no information regarding overdosage of this product.

### WHAT TO DO WHEN YOU HAVE USED MORE THAN THE RECOMMENDED DOSAGE?

If you have taken more than the recommended dosage, consult a doctor.

### CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- Follow directions for dilution and the recommended amount of this reduced osmolarity ORS solution according to weight and/or age of the patient and intended use (prevention or treatment) of dehydration to avoid underdosing or overdosing.
- Oral Rehydration Salts may be used in severe dehydration only if intravenous fluid is not available.
- This reduced osmolarity ORS granules in sachet should be made into solution immediately before use. If not refrigerated, discard any remaining solution an hour after constitution. If refrigerated, the constituted solution can be kept for 24 hours after which it should no longer be used.
- Use only cooled boiled/clean drinking water to constitute this product. The constituted solution should never be boiled.
- This product gives a cloudy solution when reconstituted. Shake or stir well before feeding.
- Do not use more than the recommended dose unless recommended by a doctor.
- Do not use after the expiry date on the label.

### WHEN SHOULD YOU CONSULT YOUR DOCTOR?

If any undesirable effect occurs or dehydration worsen.

### AVAILABILITY

Aluminum Sachet of 4.34 g (Box of 30's)

### DATE OF OF PATIENT INFORMATION LEAFLET (PIL): August 2018

FDA Registration no. DRHR-1112

### ADR Reporting Statement

"For suspected adverse drug reaction, report to the FDA: [www.fda.gov.ph](http://www.fda.gov.ph)"  
Patient must seek medical attention immediately at the first sign of any adverse drug reaction

### Manufactured by:

SAN MARINO LABORATORIES CORPORATION  
# 1 Crisanto delos Reyes St., Brgy. Javalera  
Gen. Trias, Cavite

### for:

PHARMACARE PRODUCTS CORPORATION  
SB Residences Unit 2&3 302 P. Dela Cruz St.,  
San Bartolome, Novaliches Quezon City