

## CARBOCISTEINE

### BROXYTONE®

250mg/5mL Pediatric Syrup  
Mucolytic

#### FORMULATIONS

Each 5mL (1 teaspoonful) Pediatric Syrup contains:  
Carbocisteine ..... 250mg

#### PRODUCT DESCRIPTION

Carbocisteine (Broxytone) Pediatric Syrup is a clear, green syrup with peppermint flavor.

#### WHAT IS IN THE MEDICINE?

Carbocisteine is used in the treatment of disorders of the respiratory tract with excessive mucus such as acute and chronic bronchitis, bronchiectasis, asthmatic bronchitis and emphysema. In otorhinology, it is indicated as adjunctive therapy in otitis, sinusitis, nasopharyngitis, tubal otorrhea, and clearing up hypersecretion prior surgery.

#### STRENGTH OF MEDICINE

See formulation

#### WHAT IS THE MEDICINE USED FOR?

For the treatment of disorders of the respiratory tract associated with excessive or viscous mucus.

#### HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

AGE GROUP	Broxytone 250mg / 5mL Pediatric Syrup
Children: 2 to 5 years 6 to 12 years	1.25mL -2.5mL four times a day 5mL three times daily
Or as prescribed by the physician.	

#### WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

If you are allergic to any ingredient in the product.  
Avoid Carbocisteine in patients with active stomach or intestinal ulcer.

#### CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

These medicines should be used with caution in patients with history of peptic ulcer and be avoided by patients with active ulceration.  
Do not take more than the recommended dose.  
Do not use after the expiry date on the label.

#### UNDESIRABLE EFFECTS OF THIS MEDICINE

Carbocisteine is generally safe.  
Nausea, gastric discomfort, headache, diarrhea, rashes, gastrointestinal bleeding, are most common undesirable effects.  
Carbocisteine rarely causes dizziness, insomnia, palpitations, mild lowering of blood glucose, dryness of the mouth, abnormal heart rhythm (atrial fibrillation) and "gassiness".

#### WHAT OTHER MEDICINES OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Always tell your doctor if you are taking other medicines, including herbal medicines and food supplements.

#### WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, just take the next dose if still needed for the condition being treated and the subsequent doses at the recommended time and schedule.  
Do not double the dose.

#### SIGNS AND SYMPTOMS OF OVERDOSAGE

Gastrointestinal disturbance is the most likely symptom of Carbocisteine overdose. Gastric lavage followed by observation may be beneficial therapy for overdose.

#### WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSE?

If you have taken more than the recommended dosage, consult a doctor or contact a poison control center right away.

#### WHAT SHOULD YOU CONSULT YOUR DOCTOR?

- If you are pregnant or breastfeeding, consult your doctor before using this product.
- Stop taking this medicine and consult your doctor immediately if you develop any of the following:
  - Skin rashes
  - Stomach or intestinal bleeding
- If symptoms does not improve or new symptoms appear.

#### HOW SHOULD YOU KEEP THIS MEDICINE?

Keep the product out of reach and sight of children  
Store at temperatures not exceeding 30°C

#### AVAILABILITY

250mg / 5mL Pediatric Syrup - Amber bottle x 60ml (Box of 1's)

"For suspected Adverse Drug Reaction, report to the FDA: [www.fda.gov.ph](http://www.fda.gov.ph)"

Revision date: March 01, 2018

Broxytone 250mg/5mL Pediatric Syrup: DRHR-1449

Manufactured by:  
**SAN MARINO LABORATORIES CORP.**  
No. 1 Crisanto delos Reyes Street  
Brgy. Javalera General Trias Cavite

For:  
**I . A . E**  
pharmaceuticals corp.  
#25 Kabinayan St., Quezon City