

ASCORBIC ACID

CAREZEE

100 mg/5 mL Syrup
100 mg/mL Syrup (Oral Drops)
VITAMIN

FORMULATION:

Each 5 mL contains:

Ascorbic acid, USP.....100 mg

Each mL syrup (oral drops) contains:

Ascorbic acid, USP (as Sodium Ascorbate).....112 mg).....100 mg

PRODUCT DESCRIPTION:

Carezee syrup is a clear, orange to brown orange syrup with lemon odor and flavor.

Carezee drops is a clear, orange to brown orange syrup has sweet, slightly salty taste with lemon flavor and odor.

WHAT IS IN THE MEDICINE?

This product contains Vitamin C. Vitamin C helps the body's natural defense against damaging free radicals. Free radicals are highly reactive and unstable chemicals generated during normal body activities that require oxygen and after exposure to UV light, cigarette smoke and various pollutants. Vitamin C forms part of the body's natural immune system and stimulates the activity of specialized white blood cells which inhibit harmful bacteria and viruses. Vitamin C also needed for the formation of the protein, collagen. It is important in the formation of bones, teeth, cartilage and skin and promotes faster healing.

STRENGTH OF THE MEDICINE

See formulation above

WHAT IS THE MEDICINE USED FOR?

For the treatment and prevention of Vitamin C deficiency. For body resistance against stress, common colds and other infections.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THE MEDICINE?

Orally, once a day or as prescribed by the physician

For Carezee syrup: Children - ¼ to 1 teaspoon (2.5mL - 5mL)

Adults - 2 to 4 teaspoonfuls (10mL - 20mL)

For Carezee drops: Infants - 0.6mL (60mg) daily

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

If you are allergic to any ingredient of the product.

UNDESIRABLE EFFECTS

Prolonged intake of Vitamin C in excess of 2 g/day may lead to nausea, abdominal cramps, diarrhea and nose bleeds.

WHAT OTHER MEDICINE OR FOODS SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

The amount of Vitamin C in the product is not expected to interact with food or medicines.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, just give the next dose and the subsequent doses at the usual recommended schedule. Do not double the dose unless recommended by a doctor.

HOW SHOULD YOU KEEP THE MEDICINE?

- Keep the product out of reach and sight of children.
- Store at temperatures not exceeding 30°C
- Protect from light
- Always keep container tightly closed.

SIGNS AND SYMPTOMS OF OVERDOSE

There is no information regarding overdosage of Vitamin C.

WHAT TO DO WHEN YOU HAVE USED MORE THAN THE RECOMMENDED DOSAGE?

If you have taken more than the recommended dosage, consult a doctor.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- Should be given with care to patients with hyperoxaluria.
- Do not use after the expiry date on the label.
- Do not give more than the recommended dose.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

If any undesirable effect occurs.

AVAILABILITY

Syrup: Boston round amber glass bottle x 120mL (Box of 1's)

Drops: Amber Bottle x 30 mL with Aluminum Cap, Medicine Dropper.

DATE OF REVISION OF PATIENT INFORMATION LEAFLET (PIL): February 12, 2018

Carezee Oral Drops: Registration no. DR-XY35220 Valid until December 10, 2021

Carezee Syrup: Registration no. DRHR-881 Valid until October 8, 2019.

ADR Reporting Statement

"For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph"

Patient must seek medical attention immediately at the first sign of any adverse drug reaction.

Manufactured for:

PHARMACARE PRODUCTS CORPORATION
SB Residences Unit 2&3 302 P. Dela Cruz St.,
San Bartolome, Novaliches Quezon City

Manufactured by:

LA CROSEUS PHARMA INC.
Brgy. Sto. Tomas Biñan, Laguna