EMTRICITABINE
TENOFOVIR DISOPROXIL FUMARATE

TENOF - EM

200 mg/300 mg Film-coated tablet

Antiretroviral

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE TABLETS safely and effectively. See full prescribing information for EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE TABLETS.

EMTRICITABINE and TENOFOVIR DISOPROXIL FUMARATE tablets, for oral use

Initial U.S. Approval: 2004

WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS, POST-TREATMENT ACUTE EXACERBATION OF HEPATITIS B, and RISK OF DRUG RESISTANCE WITH USE OF EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE TABLET FOR PrEP IN UNDIAGNOSED HIV-1 INFECTION

See full prescribing information for complete boxed warning.

- **Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogs, including tenofovir disoproxil fumarate, a component of emtricitabine and tenofovir disoproxil fumarate tablet. (5.1)**
- **Emtricitabine and tenofovir disoproxil fumarate is not approved for the treatment of chronic hepatitis B virus (HBV) infection. Severe acute exacerbations of hepatitis B have been reported in patients coinfected with HIV-1 and HBV who have discontinued emtricitabine and tenofovir disoproxil fumarate tablets. Therefore, hepatic function should be monitored closely in HBV-infected patients who discontinue emtricitabine and tenofovir disoproxil fumarate tablets. If appropriate, initiation of anti-hepatitis B therapy may be warranted. (5.2)**
- **Emtricitabine and tenofovir disoproxil fumarate tablets used for a PrEP indication must only be prescribed to individuals confirmed to be HIV-negative immediately prior to initial use and periodically during use. Drug-resistant HIV-1 variants have been identified with the use of emtricitabine and tenofovir disoproxil fumarate tablets for a PrEP indication following undetected acute HIV-1 infection. Do not initiate emtricitabine and tenofovir disoproxil fumarate tablets for a PrEP indication if signs or symptoms of acute HIV infection are present unless negative infection status is confirmed. (5.9)**

----------RECENT MAJOR CHANGES----------

Indications and Usage	
Treatment of HIV-1 infection (1.1)	03/2016
Dosage and Administration (2.2)	03/2016
Warnings and Precautions	
Coadministration with Other Products (5.4)	03/2016

----------INDICATIONS AND USAGE----------

Emtricitabine and tenofovir disoproxil fumarate tablet is a combination of emtricitabine and tenofovir disoproxil fumarate, both nucleoside analog HIV-1 reverse transcriptase inhibitors.

Emtricitabine and tenofovir disoproxil fumarate tablet is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and pediatric patients weighing at least 17 kg. (1)

Emtricitabine and tenofovir disoproxil fumarate tablet is indicated in combination with safer sex practices for pre-exposure prophylaxis (PrEP) to reduce the risk of sexually acquired HIV-1 in adults at high risk. (1)

----------DOSAGE AND ADMINISTRATION----------

Treatment of HIV-1 Infection (2.1)

- Recommended dose in adults and pediatric patients weighing greater than or equal to 35 kg: One emtricitabine and tenofovir disoproxil fumarate tablet (containing 200 mg/300 mg of emtricitabine and tenofovir disoproxil fumarate) once daily taken orally with or without food. (2.1)
- Recommended dose in pediatric patients weighing greater than or equal to 17 kg and able to swallow a whole tablet: one emtricitabine and tenofovir disoproxil fumarate low strength tablet (100 mg/150 mg, 133 mg/ 200 mg, or 167 mg/250 mg based on body weight) once daily taken orally with or without food. (2.2)
- Recommended dose in renally impaired HIV-1 infected adult patients: Creatinine clearance 30 to 49 mL/min: 1 tablet every 48 hours. (2.4) CrCl below 30 mL/min or hemodialysis: Do not use emtricitabine and tenofovir disoproxil fumarate tablets. (2.4)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNINGS: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS, POST TREATMENT ACUTE EXACERBATION OF HEPATITIS B, and RISK OF DRUG RESISTANCE WITH USE OF EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE FOR PRE-EXPOSURE PROPHYLAXIS (PrEP) IN UNDIAGNOSED EARLY HIV-1 INFECTION
1 INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 Pre-Exposure Prophylaxis
2 DOSAGE AND ADMINISTRATION
2.1 Recommended Dose for Treatment of HIV-1 Infection in Adults and Pediatric Patients Weighing 35 kg or More
2.2 Recommended Dose for Treatment of HIV-1 Infection in Pediatric Patients Weighing 17kg or More and Able to Swallow a Whole Tablet
2.3 Recommended Dose for Pre-exposure Prophylaxis
2.4 Dose Adjustment for Renal Impairment
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Lactic Acidosis/Severe Hepatomegaly with Steatosis
5.2 HBV Infection
5.3 New Onset or Worsening Renal Impairment
5.4 Coadministration with Other Products
5.5 Bone Effects of Tenofovir DF
5.6 Fat Redistribution
5.7 Immune Reconstitution Syndrome
5.8 Early Virologic Failure
5.9 Comprehensive Management to Reduce the Risk of Acquiring HIV-1
6 ADVERSE REACTIONS
6.1 Adverse Reactions from Clinical Trials Experience in HIV-1 Infected Subjects
6.2 Adverse Reactions from Clinical Trial Experience in HIV-1 Uninfected Adult Subjects
6.3 Postmarketing Experience
7 DRUG INTERACTIONS
7.1 Didanosine
7.2 HIV-1 Protease Inhibitors
7.3 Hepatitis C Antiviral Agents
7.4 Drugs Affecting Renal Function
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.3 Nursing Mothers
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Patients with Impaired Renal Function
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.3 Pharmacokinetics
12.4 Microbiology
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology and/or Pharmacology
14 CLINICAL STUDIES
14.1 Study 934
14.2 iPrEx Trial
14.3 Partners PrEP Trial
16 HOW SUPPLIED/STORAGE AND HANDLING
17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

WARNING: LACTIC ACIDOSIS/SEVERE HEPATOMEGALY WITH STEATOSIS, POST TREATMENT ACUTE EXACERBATION OF HEPATITIS B, and RISK OF DRUG RESISTANCE WITH USE OF EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE TABLETS FOR PRE-EXPOSURE PROPHYLAXIS (PrEP) IN UNDIAGNOSED EARLY HIV-1 INFECTION

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogs, including tenofovir disoproxil fumarate, a component of emtricitabine and tenofovir disoproxil fumarate tablets, in combination with other antiretrovirals *(See Warnings and Precautions (5.1))*

Emtricitabine and tenofovir disoproxil fumarate is not approved for the treatment of chronic hepatitis B virus (HBV) infection and the safety and efficacy of emtricitabine and tenofovir disoproxil fumarate tablets have not been established in patients coinfected with HBV and HIV-1. Severe acute exacerbations of hepatitis B have been reported in patients who are coinfected with HBV and HIV-1 and have discontinued emtricitabine and tenofovir disoproxil fumarate tablets. Therefore, hepatic function should be monitored closely with both clinical and laboratory follow-up for at least several months in patients who are infected with HBV and discontinue emtricitabine and tenofovir disoproxil fumarate. If appropriate, initiation of anti-hepatitis B therapy may be warranted *(See Warnings and Precautions (5.2))*.

Emtricitabine and tenofovir disoproxil fumarate tablets used for a PrEP indication must only be prescribed to individuals confirmed to be HIV-negative immediately prior to initiating and periodically (at least every 3 months) during use. Drug-resistant HIV-1 variants have been identified with use of emtricitabine and tenofovir disoproxil fumarate tablets for a PrEP indication following undetected acute HIV-1 infection. Do not initiate emtricitabine and tenofovir disoproxil fumarate tablets for a PrEP indication if signs or symptoms of acute HIV-1 infection are present unless negative infection status is confirmed *(See Warnings and Precautions (5.9))*.

1 INDICATIONS AND USAGE

1.1 Treatment of HIV-1 Infection

Emtricitabine and tenofovir disoproxil fumarate tablet, a combination of emtricitabine and tenofovir disoproxil fumarate, is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and pediatric patients weighing at least 17 kg. *[see Dosage and Administration (2) and Clinical Studies (14)]*.

The following points should be considered when initiating therapy with emtricitabine and tenofovir disoproxil fumarate tablets for the treatment of HIV-1 infection:

- It is not recommended that emtricitabine and tenofovir disoproxil fumarate tablet be used as a component of a triple nucleoside regimen.
- Emtricitabine and tenofovir disoproxil fumarate tablet should not be coadministered with ATRIPLA®, COMPLERA®, emtricitabine, GENVOYA®, ODEFSEY®, STRIBILD®, tenofovir disoproxil fumarate or lamivudine-containing products *[see Warnings and Precautions (5.4)]*.
- In treatment experienced patients, the use of emtricitabine and tenofovir disoproxil fumarate tablet should be guided by laboratory testing and treatment history *[See Microbiology (12.4)]*.

1.2 Pre-Exposure Prophylaxis

Emtricitabine and tenofovir disoproxil fumarate tablet is indicated in combination with safer sex practices for pre-exposure prophylaxis (PrEP) to reduce the risk of sexually acquired HIV-1 in adults at high risk. This indication is based on clinical trials in men who have sex with men (MSM) at high risk for HIV-1 infection and in heterosexual serodiscordant couples *[See Clinical Studies (14.2, 14.3)]*.

When considering emtricitabine and tenofovir disoproxil fumarate tablet for pre-exposure prophylaxis the following factors may help to identify individuals at high risk:

- has partner(s) known to be HIV-1 infected, or
- engages in sexual activity within a high prevalence area or social network and one or more of the following:
  - inconsistent or no condom use
  - diagnosis of sexually transmitted infections
  - exchange of sex for commodities (such as money, food, shelter, or drugs)
  - use of illicit drugs or alcohol dependence
  - incarceration
  - partner(s) of unknown HIV-1 status with any of the factors listed above

When prescribing emtricitabine and tenofovir disoproxil fumarate tablet for pre-exposure prophylaxis, healthcare providers must:

- prescribe emtricitabine and tenofovir disoproxil fumarate tablet as part of a comprehensive prevention strategy because emtricitabine and tenofovir disoproxil fumarate tablet is not always effective in preventing the acquisition of HIV-1 infection *[See Warnings and Precautions (5.9)]*;
- counsel all uninfected individuals to strictly adhere to the recommended emtricitabine and tenofovir disoproxil fumarate tablet dosing schedule because the effectiveness of emtricitabine and tenofovir disoproxil fumarate tablet in reducing the risk of acquiring HIV-1 was strongly correlated with adherence as demonstrated by measurable drug levels in clinical trials *[See Warnings and Precautions (5.9)]*;
- confirm a negative HIV-1 test immediately prior to initiating emtricitabine and tenofovir disoproxil fumarate tablet for a PrEP indication. If clinical symptoms consistent with acute viral infection are present and recent (<1 month) exposures are suspected, delay starting PrEP for at least one month and reconfirm HIV-1 status or use a test approved by the FDA as an aid in the diagnosis of HIV-1 infection, including acute or primary HIV-1 infection. *[See Warnings and Precautions (5.9)]*; and
- screen for HIV-1 infection at least once every 3 months while taking emtricitabine and tenofovir disoproxil fumarate tablet for PrEP.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose for Treatment of HIV-1 Infection in Adults and Pediatric Patients Weighing 35 Kg or More

The recommended dose of emtricitabine and tenofovir disoproxil fumarate tablets in adults and in pediatric patients with body weight greater than or equal to 35 kg is one tablet (containing 200 mg of emtricitabine and 300 mg of tenofovir disoproxil fumarate) once daily taken orally with or without food.

2.2 Recommended Dose for Treatment of HIV-1 Infection in Pediatric Patients Weighing at Least 17kg or More and Able to Swallow a Whole Tablet

The recommended oral dose for pediatric patients weighing greater than or equal to 17 kg and who are able to swallow a whole tablet, is one emtricitabine and tenofovir disoproxil fumarate low strength tablet (emtricitabine [FTC]/tenofovir disoproxil fumarate [TDF]) (167 mg/250 mg, 133 mg/200 mg, or 100 mg/150 mg based on body weight) taken orally once daily with or without food.

The recommended oral dosage of emtricitabine and tenofovir disoproxil fumarate low strength tablets is presented in Table 1. Weight should be monitored periodically and the emtricitabine and tenofovir disoproxil fumarate tablet dose adjusted accordingly.

Table 1 Dosing for Pediatric Patients Weighing 17 kg to less than 35 kg using Emtricitabine and Tenofovir disoproxil fumarate Low Strength Tablets

Body Weight (kg)	Dosing of FTC (mg)/TDF (mg)
17 to less than 22	one 100/150 tablet once daily
22 to less than 28	one 133/200 tablet once daily
28 to less than 35	one 167/250 tablet once daily

2.3 Recommended Dose for Pre-exposure Prophylaxis

The dose of emtricitabine and tenofovir disoproxil fumarate tablet in HIV-1 uninfected adults is one tablet (containing 200 mg of emtricitabine and 300 mg of tenofovir disoproxil fumarate) once daily taken orally with or without food.

2.4 Dose Adjustment for Renal Impairment

Treatment of HIV-1 Infection

Significantly increased drug exposures occurred when emtricitabine or tenofovir disoproxil fumarate tablets were administered to subjects with moderate to severe renal impairment *[See emtricitabine or tenofovir disoproxil fumarate Package Insert]*. Therefore, adjust the dosing interval of emtricitabine and tenofovir disoproxil fumarate tablet in HIV-1 infected adult patients with baseline creatinine clearance 30 to 49 mL/min using the recommendations in Table 2. These dosing interval recommendations are based on modeling of single-dose pharmacokinetic data in non-HIV infected subjects. The safety and effectiveness of these dosing interval adjustment recommendations have not been clinically evaluated in patients with moderate renal impairment, therefore clinical response to treatment and renal function should be closely monitored in these patients *[See Warnings and Precautions (5.3)]*.

No dose adjustment is necessary for HIV-1 infected patients with mild renal impairment (creatinine clearance 50 to 80 mL/min). No data are available to make dose recommendations in pediatric patients with renal impairment.

Table 2 Dosage Adjustment for HIV-1 Infected Adult Patients with Altered Creatinine Clearance

	Creatinine Clearance (mL/min)[a]		
<30	**≥50**	**30–49**	**(Including Patients Requiring Hemodialysis)**
Recommended Dosing Interval	Every 24 hours	Every 48 hours	Emtricitabine and Tenofovir disoproxil fumarate should not be administered.

a. Calculated using ideal (lean) body weight

Routine monitoring of estimated creatinine clearance, serum phosphorus, urine glucose, and urine protein should be performed in all individuals with mild renal impairment *[See Warnings and Precautions (5.3)]*.

Pre-exposure Prophylaxis

Do not use emtricitabine and tenofovir disoproxil fumarate tablet for a PrEP indication in HIV-1 uninfected individuals with estimated creatinine clearance below 60 mL/min *[See Warnings and Precautions (5.3)]*.

Routine monitoring of estimated creatinine clearance, serum phosphorus, urine glucose, and urine protein should be performed in all individuals with mild renal impairment. If a decrease in estimated creatinine clearance is observed in uninfected individuals while using emtricitabine and tenofovir disoproxil fumarate tablet for PrEP, evaluate potential causes and re-assess potential risks and benefits of continued use *[See Warnings and Precautions (5.3)]*.

3 DOSAGE FORMS AND STRENGTHS

Emtricitabine and Tenofovir disoproxil fumarate is available as tablets. Each tablet contains 200 mg of emtricitabine and 300 mg of tenofovir disoproxil fumarate (which is equivalent to 245 mg of tenofovir disoproxil). The tablets are blue, capsule shaped, film coated tablets, debossed with 'H' on one side and with '124' on the other side.

4 CONTRAINDICATIONS

Do not use emtricitabine and tenofovir disoproxil fumarate tablets for pre-exposure prophylaxis in individuals with unknown or positive HIV-1 status. Emtricitabine and tenofovir disoproxil fumarate tablets should be used in HIV-infected patients only in combination with other antiretroviral agents

5 WARNINGS AND PRECAUTIONS

5.1 Lactic Acidosis/Severe Hepatomegaly with Steatosis

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogs, including tenofovir disoproxil fumarate, a component of emtricitabine and tenofovir disoproxil fumarate tablet, in combination with other antiretrovirals. A majority of these cases have been in women. Obesity and prolonged nucleoside exposure may be risk factors. Particular caution should be exercised when administering nucleoside analogs to any patient or uninfected individual with known risk factors for liver disease, however, cases have also been reported in HIV-1 infected patients with no known risk factors. Treatment with emtricitabine and tenofovir disoproxil fumarate should be suspended in any patient or uninfected individual who develops clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity (which may include hepatomegaly and steatosis even in the absence of marked transaminase elevations).

5.2 HBV Infection

It is recommended that all individuals be tested for the presence of chronic hepatitis B virus (HBV) before initiating emtricitabine and tenofovir disoproxil fumarate. Emtricitabine and Tenofovir disoproxil fumarate is not approved for the treatment of chronic HBV infection and the safety and efficacy of emtricitabine and tenofovir disoproxil fumarate have not been established in patients infected with HBV. Severe acute exacerbations of hepatitis B have been reported in patients who are coinfected with HBV and HIV-1 and have discontinued emtricitabine and tenofovir disoproxil fumarate. In some patients infected with HBV and treated with emtricitabine, the exacerbations of hepatitis B were associated with liver decompensation and liver failure. Patients who are infected with HBV should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment with emtricitabine and tenofovir disoproxil fumarate. If appropriate, initiation of anti-hepatitis B therapy may be warranted. HBV-uninfected individuals should be offered vaccination.

5.3 New Onset or Worsening Renal Impairment

Emtricitabine and tenofovir are principally eliminated by the kidney. Renal impairment, including cases of acute renal failure and Fanconi syndrome (renal tubular injury with severe hypophosphatemia), has been reported with the use of tenofovir disoproxil fumarate *[See Adverse Reactions (6.3)]*.

It is recommended that estimated creatinine clearance be assessed in all individuals prior to initiating therapy and as clinically appropriate during therapy with emtricitabine and tenofovir disoproxil fumarate. In patients at risk of renal dysfunction, including patients who have previously experienced renal events while receiving HEPSERA®, it is recommended that estimated creatinine clearance, serum phosphorus, urine glucose, and urine protein be assessed prior to initiation of emtricitabine and tenofovir disoproxil fumarate, and periodically during emtricitabine and tenofovir disoproxil fumarate therapy.

Emtricitabine and tenofovir disoproxil fumarate should be avoided with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple non-steroidal anti-inflammatory drugs (NSAIDs)) *[See Drug Interactions (7.4)]*.

Cases of acute renal failure after initiation of high dose or multiple NSAIDs have been reported in HIV-infected patients with risk factors for renal dysfunction who appeared stable on tenofovir disoproxil fumarate (tenofovir DF). Some patients required hospitalization and renal replacement therapy. Alternatives to NSAIDs should be considered, if needed, in patients at risk for renal dysfunction.

Persistent or worsening bone pain, pain in extremities, fractures and/or muscular pain or weakness may be manifestations of proximal renal tubulopathy and should prompt an evaluation of renal function in at-risk patients.

Treatment of HIV-1 Infection

Dosing interval adjustment of emtricitabine and tenofovir disoproxil fumarate and close monitoring of renal function are recommended in all patients with estimated creatinine clearance 30 to 49 mL/min *[See Dosage and Administration (2.4)]*. No safety or efficacy data are available in patients with renal impairment who received emtricitabine and tenofovir disoproxil fumarate using these dosing guidelines, so the potential benefit of emtricitabine and tenofovir disoproxil fumarate therapy should be assessed against the potential risk of renal toxicity. Emtricitabine and Tenofovir disoproxil fumarate should not be administered to patients with estimated creatinine clearance below 30 mL/min or patients requiring hemodialysis.

Pre-exposure Prophylaxis

Emtricitabine and Tenofovir disoproxil fumarate for a PrEP indication should not be used if estimated creatinine clearance is less than 60 mL/min. If a decrease in estimated creatinine clearance is observed in uninfected individuals while using emtricitabine and tenofovir disoproxil fumarate for PrEP, evaluate potential causes and re-assess potential risks and benefits of continued use *[See Dosage and Administration (2.4)]*.

5.4 Coadministration with Other Products

Emtricitabine and tenofovir disoproxil fumarate is a fixed-dose combination of emtricitabine and tenofovir disoproxil fumarate. Do not coadminister emtricitabine and tenofovir disoproxil fumarate with other drugs containing emtricitabine or tenofovir disoproxil fumarate, or containing tenofovir alafenamide, including ATRIPLA, COMPLERA, emtricitabine, GENVOYA, ODEFSEY, STRIBILD, or tenofovir disoproxil fumarate. Due to similarities between emtricitabine and lamivudine, do not coadminister emtricitabine and tenofovir disoproxil fumarate with other drugs containing lamivudine, including Combivir (lamivudine/zidovudine), Dutrebis (lamivudine/raltegravir), Epivir or Epivir-HBV (lamivudine), Epzicom (abacavir sulfate/lamivudine), Triumeq (abacavir sulfate/dolutegravir/lamivudine), or Trizivir (abacavir sulfate/lamivudine/zidovudine).

Do not coadminister emtricitabine and tenofovir disoproxil fumarate with HEPSERA (adefovir dipivoxil).

5.5 Bone Effects of Tenofovir DF

Bone Mineral Density :

In clinical trials in HIV-1 infected adults and in a clinical trial of HIV-1 uninfected individuals, tenofovir DF was associated with slightly greater decreases in bone mineral density (BMD) and increases in biochemical markers of bone metabolism, suggesting increased bone turnover relative to comparators. *[See Adverse Reactions (6.2)* and tenofovir disoproxil fumarate prescribing information]. Serum parathyroid hormone levels and 1,25 Vitamin D levels were also higher in subjects receiving tenofovir] DF.

Clinical trials evaluating tenofovir DF in pediatric and adolescent subjects were conducted. Under normal circumstances, BMD increases rapidly in pediatric patients. In HIV-1 infected subjects aged 2 years to less than 18 years, bone effects were similar to those observed in adult subjects and suggest increased bone turnover. Total body BMD gain was less in the tenofovir DF treated HIV-1 infected pediatric subjects as compared to the control groups. Similar trends were observed in chronic hepatitis B infected adolescent subjects aged 12 years to less than 18 years. In all pediatric trials, skeletal growth (height) appeared to be unaffected. For more information, please consult the tenofovir disoproxil fumarate prescribing information.

The effects of tenofovir DF-associated changes in BMD and biochemical markers on long-term bone health and future fracture risk are unknown. Assessment of BMD should be considered for adult and pediatric patients who have a history of pathologic bone fracture or other risk factors for osteoporotic bone loss. Although the effect of supplementation with calcium and vitamin D was not studied, such supplementation may be beneficial. If bone abnormalities are suspected then appropriate consultation should be obtained.

Mineralization Defects:

Cases of osteomalacia associated with proximal renal tubulopathy, manifested as bone pain or pain in extremities and which may contribute to fractures, have been reported in association with the use of tenofovir DF *[See Adverse Reactions (6.3)]*. Arthralgias and muscle pain or weakness have also been reported in cases of proximal renal tubulopathy. Hypophosphatemia and osteomalacia secondary to proximal renal tubulopathy should be considered in patients at risk of renal dysfunction who present with persistent or worsening bone or muscle symptoms while receiving products containing tenofovir DF *[See Warnings and Precautions (5.3)]*.

5.6 Fat Redistribution

Redistribution/accumulation of body fat including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and "cushingoid appearance" have been observed in HIV-1 infected patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

5.7 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in HIV-1 infected patients treated with combination antiretroviral therapy, including emtricitabine and tenofovir disoproxil fumarate. During the initial phase of combination antiretroviral treatment, HIV-1 infected patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium infection, cytomegalovirus, Pneumocystis jirovecii pneumonia* (PCP), or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, and Guillain-Barré syndrome) have also been reported to occur in the setting of immune reconstitution, however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.8 Early Virologic Failure

Clinical trials in HIV-1 infected subjects have demonstrated that certain regimens that only contain three nucleoside reverse transcriptase inhibitors (NRTI) are generally less effective than triple drug regimens containing two NRTIs in combination with either a non-nucleoside reverse transcriptase inhibitor or a HIV-1 protease inhibitor. In particular, early virological failure and high rates of resistance substitutions have been reported. Triple nucleoside regimens should therefore be used with caution. Patients on a therapy utilizing a triple nucleoside-only regimen should be carefully monitored and considered for treatment modification.

5.9 Comprehensive Management to Reduce the Risk of Acquiring HIV-1

Use emtricitabine and tenofovir disoproxil fumarate for pre-exposure prophylaxis only as part of a comprehensive prevention strategy that includes other prevention measures, such as safer sex practices, because emtricitabine and tenofovir disoproxil fumarate is not always effective in preventing the acquisition of HIV-1 *[See Clinical Studies (14.2 and 14.3)]*.

- Counsel uninfected individuals about safer sex practices that include consistent and correct use of condoms, knowledge of their HIV-1 status and that of their partner(s), and regular testing for other sexually transmitted infections that can facilitate HIV-1 transmission (such as syphilis and gonorrhea).
- Inform uninfected individuals about and support their efforts in reducing sexual risk behavior.

Pre-exposure Prophylaxis (2.3)

- Recommended dose in HIV-1 uninfected adults: One tablet (containing 200 mg/300 mg of emtricitabine and tenofovir disoproxil fumarate) once daily taken orally with or without food. (2.3)
- Recommended dose in renally impaired HIV-uninfected individuals: Do not use emtricitabine and tenofovir disoproxil fumarate tablets in HIV-uninfected individuals if CrCl is below 60 mL/min. If a decrease in CrCl is observed in uninfected individuals while using emtricitabine and tenofovir disoproxil fumarate tablets for PrEP, evaluate potential causes and re-assess potential risks and benefits of continued use. (2.4)

----------DOSAGE FORMS AND STRENGTHS----------

Tablets: 200 mg of emtricitabine and 300 mg of tenofovir disoproxil fumarate. (3)

----------CONTRAINDICATIONS----------

Do not use emtricitabine and tenofovir disoproxil fumarate tablets for pre-exposure prophylaxis in individuals with unknown or positive HIV-1 status. Emtricitabine and tenofovir disoproxil fumarate tablets should be used in HIV-infected patients only in combination with other antiretroviral agents. (4)

----------WARNINGS AND PRECAUTIONS----------

- New onset or worsening renal impairment: Can include acute renal failure and Fanconi syndrome. Assess estimated creatinine clearance before initiating treatment with emtricitabine and tenofovir disoproxil fumarate. In patients at risk for renal dysfunction, assess estimated creatinine clearance, serum phosphorus, urine glucose and urine protein before initiating treatment with emtricitabine and tenofovir disoproxil fumarate and periodically during treatment. Avoid administering emtricitabine and tenofovir disoproxil fumarate with concurrent or recent use of nephrotoxic drugs. (5.3)
- Coadministration with Other Products: Do not use with drugs containing emtricitabine, tenofovir alafenamide or tenofovir disoproxil fumarate including ATRIPLA, COMPLERA, emtricitabine, GENVOYA, ODEFSEY, STRIBILD, tenofovir disoproxil fumarate ; or with drugs containing lamivudine. Do not administer in combination with HEPSERA. (5.4)
- Decreases in bone mineral density (BMD): Consider assessment of BMD in patients with a history of pathologic fracture or other risk factors for osteoporosis or bone loss. (5.5)
- Redistribution/accumulation of body fat: Observed in patients receiving antiretroviral therapy. (5.6)
- Immune reconstitution syndrome: May necessitate further evaluation and treatment. (5.7)
- Triple nucleoside-only regimens: Early virologic failure has been reported in HIV-infected patients. Monitor carefully and consider treatment modification. (5.8)
- Comprehensive management to reduce the risk of acquiring HIV-1: Use as part of a comprehensive prevention strategy including other prevention measures; strictly adhere to dosing schedule. (5.9)
- Management to reduce the risk of acquiring HIV-1 drug resistance:
  Prior to initiating emtricitabine and tenofovir disoproxil fumarate for PrEP -if clinical symptoms consistent with acute viral infection are present and recent (<1 month) exposures are suspected, delay starting PrEP for at least one month and reconfirm negative HIV-1 status or use a test approved by the FDA as an aid in the diagnosis of HIV-1 infection, including acute or primary HIV-1 infection.
  While using emtricitabine and tenofovir disoproxil fumarate for PrEP - HIV-1 screening tests should be repeated at least every 3 months. (5.9)

----------ADVERSE REACTIONS----------

In HIV-1 infected patients, the most common adverse reactions (incidence greater than or equal to 10%) are diarrhea, nausea, fatigue, headache, dizziness, depression, insomnia, abnormal dreams, and rash. (6.1)

In HIV-1 uninfected individuals in PrEP trials, adverse reactions that were reported by more than 2% of emtricitabine and tenofovir disoproxil fumarate subjects and more frequently than by placebo subjects were headache, abdominal pain and weight decreased. (6.2)

----------DRUG INTERACTIONS----------

- Didanosine: Tenofovir disoproxil fumarate increases didanosine concentrations. Use with caution and monitor for evidence of didanosine toxicity (e.g., pancreatitis, neuropathy) when coadministered. Consider dose reductions or discontinuations of didanosine if warranted. (7.1)
- HIV-1 protease inhibitors: Coadministration decreases atazanavir concentrations and increases tenofovir concentrations. When coadministered with emtricitabine and tenofovir disoproxil fumarate, use atazanavir given with ritonavir. Coadministration of emtricitabine and tenofovir disoproxil fumarate with atazanavir and ritonavir, darunavir and ritonavir, or lopinavir/ritonavir increases tenofovir concentrations. Monitor for evidence of tenofovir toxicity. (7.2)

----------USE IN SPECIFIC POPULATIONS----------

- Nursing mothers: Women infected with HIV-1 should be instructed not to breast feed. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 05/2016

7 DRUG INTERACTIONS

No drug interaction trials have been conducted using emtricitabine and tenofovir disoproxil fumarate tablets. Drug interaction trials have been conducted with emtricitabine and tenofovir disoproxil fumarate, the components of emtricitabine and tenofovir disoproxil fumarate tablet. This section describes clinically relevant drug interactions observed with emtricitabine and tenofovir disoproxil fumarate *[see Clinical Pharmacology (12.3)]*.

7.1 Didanosine

Coadministration of emtricitabine and tenofovir disoproxil fumarate and didanosine should be undertaken with caution and patients receiving this combination should be monitored closely for didanosine-associated adverse reactions. Didanosine should be discontinued in patients who develop didanosine-associated adverse reactions.

When tenofovir disoproxil fumarate was administered with didanosine the C_{max} and AUC of didanosine increased significantly *[See Clinical Pharmacology (12.3)]*. The mechanism of this interaction is unknown. Higher didanosine concentrations could potentiate didanosine-associated adverse reactions, including pancreatitis, and neuropathy. Suppression of CD4+ cell counts has been observed in patients receiving tenofovir DF with didanosine 400 mg daily.

In patients weighing greater than 60 kg, the didanosine dose should be reduced to 250 mg when it is coadministered with emtricitabine and tenofovir disoproxil fumarate. Data are not available to recommend a dose adjustment of didanosine for adult or pediatric patients weighing less than 60 kg. When coadministered, emtricitabine and tenofovir disoproxil fumarate and Videx EC may be taken under fasted conditions or with a light meal (less than 400 kcal, 20% fat).

7.2 HIV-1 Protease Inhibitors

Tenofovir decreases the AUC and C_{min} of atazanavir *[See Clinical Pharmacology (12.3)]*. When coadministered with emtricitabine and tenofovir disoproxil fumarate, it is recommended that atazanavir 300 mg is given with ritonavir 100 mg. Emtricitabine and tenofovir disoproxil fumarate should not be coadministered with atazanavir without ritonavir.

Lopinavir/ritonavir, atazanavir coadministered with ritonavir, and darunavir coadministered with ritonavir have been shown to increase tenofovir concentrations *[See Clinical Pharmacology (12.3)]*. Tenofovir disoproxil fumarate is a substrate of P-glycoprotein (Pgp) and breast cancer resistance protein (BCRP) transporters. When tenofovir disoproxil fumarate is co-administered with an inhibitor of these transporters, an increase in absorption may be observed. Patients receiving emtricitabine and tenofovir disoproxil fumarate concomitantly with lopinavir/ritonavir, ritonavir-boosted atazanavir, or ritonavir-boosted darunavir should be monitored for tenofovir disoproxil fumarate -associated adverse reactions. Emtricitabine and Tenofovir disoproxil fumarate should be discontinued in patients who develop tenofovir disoproxil fumarate-associated adverse reactions.

7.3 Hepatitis C Antiviral Agents

Coadministration of tenofovir disoproxil fumarate and HARVONI® (ledipasvir/sofosbuvir) has been shown to increase tenofovir exposure *[See Clinical Pharmacology (12.3)]*.

In patients receiving emtricitabine and tenofovir disoproxil fumarate concomitantly with HARVONI and an HIV-1 protease inhibitor/ritonavir or an HIV-1 protease inhibitor/cobicistat combination, monitor for adverse reactions associated with tenofovir disoproxil fumarate.

In patients receiving emtricitabine and tenofovir disoproxil fumarate concomitantly with HARVONI and an HIV-1 protease inhibitor/ritonavir or an HIV-1 protease inhibitor/cobicistat combination, consider an alternative HCV or antiretroviral therapy, as the safety of increased tenofovir concentrations in this setting has not been established. If coadministration is necessary, monitor for adverse reactions associated with tenofovir disoproxil fumarate.

7.4 Drugs Affecting Renal Function

Emtricitabine and tenofovir are primarily excreted by the kidneys by a combination of glomerular filtration and active tubular secretion *[See Clinical Pharmacology (12.3)]*. No drug-drug interactions due to competition for renal excretion have been observed; however, coadministration of emtricitabine and tenofovir disoproxil fumarate with drugs that are eliminated by active tubular secretion may increase concentrations of emtricitabine, tenofovir, and/or the coadministered drug. Some examples include, but are not limited to acyclovir, adefovir dipivoxil, cidofovir, ganciclovir, valacyclovir, valganciclovir, aminoglycosides (e.g., gentamicin), and high-dose or multiple NSAIDs *[See Warnings and Precautions (5.3)]*. Drugs that decrease renal function may increase concentrations of emtricitabine and/or tenofovir.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy: *Teratogenic Effects:*

Pregnancy Category B

Antiretroviral Pregnancy Registry: To monitor fetal outcomes of pregnant women exposed to emtricitabine and tenofovir disoproxil fumarate, an Antiretroviral Pregnancy Registry (APR) has been established.

Risk Summary

Emtricitabine and Tenofovir disoproxil fumarate has been evaluated in a limited number of women during pregnancy and postpartum. Available human and animal data suggest that emtricitabine and tenofovir disoproxil fumarate does not increase the risk of major birth defects overall compared to the background rate. There are, however, no adequate and well-controlled trials in pregnant women. Because the studies in humans cannot rule out the possibility of harm, emtricitabine and tenofovir disoproxil fumarate should be used during pregnancy only if clearly needed. If an uninfected individual becomes pregnant while taking emtricitabine and tenofovir disoproxil fumarate for a PrEP indication, careful consideration should be given to whether use of emtricitabine and