

MULTIVITAMINS

INTERVIT

SYRUP VITAMINS

FORMULATION:

Ascorbic Acid (Vit. C)	75 mg
Niacinamide	10 mg
Thiamine Hydrochloride (Vit. B1)	5 mg
Pyridoxine Hydrochloride (Vit. B6)	2 mg
Riboflavin (Vit. B2)	2 mg
di-Alpha Tocopheryl Acetate (Vit. E)	2 mg
Retinol Palmitate (Vit. A)	1,1 mg
Cholecalciferol (Vit. D3)	10 mcg
Cyanocobalamin (Vit. B12)	1 mcg

DESCRIPTION:

INTERVIT a daily supplement syrup that provides the nutritional needs of children particularly during periods of rapid growth and increased physical activity.

INTERVIT is a clear, yellow - orange to brownish orange syrup with Lemon and Orange flavor. Available in Boston Round Amber bottle x 60 mL and 120 mL, with Aluminum ROPP Cap with FPE Liner (Box of 1's)

WHAT IS IN THE MEDICINE?

This Multivitamins (INTERVIT) contains It contains Vitamin B Complex vitamin to help optimize the conversion of food into energy that the body can utilize for numerous physiologic processes such as respiration, digestion, blood circulation, immune system response and as fuel by physical activity.

INTERVIT also contains fat soluble Vitamin A, Vitamin D, and Vitamin E essential for development of epithelial tissues and body resistance to disease, infection and other stressful conditions.

STRENGTH OF THE MEDICINE: See Formulaion

WHAT IS THIS MEDICINE USED FOR

A nutritional supplement to meet the nutritional needs of children during the period of increased physical activity, rapid growth and development.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

Infant : 2.5 mL (1/2 to 1 teaspoonful)
1 - 3 years : 2.5 - 5 mL (1/2 to 1 teaspoonful)
4 - 12 year : 5 mL (1 teaspoonful)
Adult : 15mL (1 tablespoonful)

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

If you are allergic to any component of this product.

CARE THAT SDHOULD BE TAKEN WEHEN TAKING THIS MEDICINE?

Avoid taking more than one multivitamin product at the same time unless your doctor tells to you.

UNDESIRABLE EFFECTS :

No common side effects have been reported with the proper use of this medicine.

seek medical attention right away if any of this severe side effects occur:

* Skin rashes * Hives * Itching * Difficulty in breathing
* Tightness in the chest * Swelling of the mouth, face, lips or tongue

WHAT OTHER MEDICINE OR FOODS SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Avoid taking more than one multivitamin product at the same time unless your doctor tells to you. Multivitamins can interact with certain medications, or affect how medications work to your body. Ask a doctor or pharmacist if it is safe for you to use multivitamins is you are also using any of the following drugs:

* Tretinoin or isotretinoin * Antibiotic
* Diuretic or water pill * Sulfa Drug or * NSAID's

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose for the day, just take the next dose and the subsequent doses at the usual recommended schedule e. i. once a day.

Do not double the dose unless recommended by a doctor.

SIGNS AND SYMPTOMS OF OVERDOSE:

Overdose symptoms may include stomach pain, vomiting, diarrhea, constipation, and loss of appetite, hair loss, peeling skin, tingly feeling in or around the mouth and headache,

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

Seek emergency medical attention as soon as possible. Call Poison Control Center right away.

HOW SHOULD YOU KEEP THIS MEDICINE?

Store at temperatures not exceeding 30°C

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

If any undesirable effects occur consult your doctor.

ADR Reporting Statement:

For suspected adverse drug reaction, report to the FDA [www.fda.gov.ph](http://www.fda.gov/ph).

Seek medical attention immediately at the first sign of any adverse drug reaction

AVAILABILITY:

Boston Round Amber bottle x 60 mL and 120 mL, with Aluminum ROPP Cap with FPE Liner (Box of 1's)

FDA REGISTRATION NUMBER: DRP-4626

DATE OF FIRST AUHORIZATION/INITIAL OF THE AUTHORIZATION
25 October 2018

DATE OF REVISION : 20 January 2020



Manufactured by:
SAN MARINO Laboratories, CORP.
1 Crisanto de los Reyes St.,
Brgy. Javalera, Gen. Trias, Cavite