

**CHOLECALCIFEROL
(VITAMIN D₃)**

**RAYMAX
3,000 IU
(equivalent 75 mcg)
CAPSULE
VITAMIN**

FORMULATION:

Each Capsule contains:
(Cholecalciferol)Vitamin D₃ 3,000 IU
(equivalent to 75 mcg)

PRODUCT DESCRIPTION:

Vitamin D (cholecalciferol-D₃) is a fat-soluble vitamin that helps your body absorb calcium and phosphorus. Having the right amount of vitamin D, calcium, and phosphorus is important for building and keeping strong bones. RAYMAX is available in gray (body) / green (cap) hard gelatin capsule; Blister Pack of 10's (Box of 60's and 100's)

WHAT IS IN THE MEDICINE?

Cholecalciferol(Vitamin D₃) is a fat-soluble vitamin that helps your body absorb calcium and phosphorus. Having the right amount of vitamin D, calcium, and phosphorus is important for building and keeping strong bones. Vitamin D is made by the body when skin is exposed to sunlight.

STRENGTH OF THE MEDICINE- See Formulation

WHAT IS THIS MEDICINE USED FOR?

Cholecalciferol is used to treat or prevent many conditions caused by lack of vitamin D, especially conditions of the skin or bones. Vitamin D is used to treat and prevent bone disorders (such as rickets, osteomalacia)

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

Take 1 (one) capsule daily. Or as prescribed by the physician.

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

*If you are allergic to any ingredient in the product

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE?

This medicine should be given with care in patients with a history of stomach or intestinal ulcer and gastrointestinal bleeding. Do not use after the expiry date on the label.

UNDESIRABLE EFFECTS

Too much vitamin D can cause harmful high calcium levels. Tell your doctor right away if any of these signs of high vitamin D/calcium levels occur: nausea/vomiting, constipation, loss of appetite, increased thirst, increased urination, mental/mood changes, unusual tiredness.

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Always tell your doctor if you are taking other medicine.

If you use other drugs or over the counter products at the same time, the effects of Vitamin D₃ may change. This may increase your risk for side-effects or cause your drug not to work properly.

Tell your doctor about all the drugs, vitamins, and herbal supplements you are using, so that you doctor can help you prevent or manage drug interactions. Vitamin D may interact with the following drugs and products, Actinomycin, Barbiturates, Cholestyramine, Glucocorticoids Paraffin oil and Phenytoin.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, just take the next dose if still needed for the condition being treated, and the subsequent doses at the recommended time or schedule. do not double the dose.

SIGNS AND SYMPTOMS OF OVERDOSE:

If someone has overdosed and has serious symptoms such as passing out or trouble breathing.

WHAT TO DO IF HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

Consult a doctor if you have taken more than the recommended dose. Call a poison control center right away.

HOW SHOULD YOU KEEP THIS MEDICINE?

Keep the product out of reach and sight of children
Store at room temperatures not exceeding 30°C

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

Too much Vitamin D can cause harmful high calcium levels. Tell your doctor right away if any of these signs of high vitamin D/calcium levels occur: nausea /vomiting, constipation, loss of appetite, increased thirst, increased urination, mental/mood changes, unusual tiredness.

ADR REPORTING STATEMENT:

"For suspected adverse drug reaction, report to the FDA: " www.fda.gov.ph " Patient must seek medical attention immediately at the first sign of any adverse drug reaction.

AVAILABILITY :

Alu/PVC red: Blister Pack of 10's (Box of 60's and 100's)

Shelf-life : 36 Months

DRP- 9201 - 02

Date of Revision :

Manufactured by:
SAN MARINO Laboratories CORP.
#1 Crisanto delos Reyes St. Javalera,
Gen. Trias, Cavite

Distributed by:
JOHTANN INTERNATIONALE PHARMA CORP.
25 Kabignayan St., Quezon City