

Vitex negundo L.
(Lagundi leaf)

NEGACOF

300 mg/5 mL Syrup
Anti-cough / Antiasthma

FORMULATION:

Each 5 mL of syrup contains the extract from 300 mg of Lagundi (*Vitex negundo L. Fam Verbenaceae*), PP dried leaves.

PRODUCT DESCRIPTION:

Negacof syrup is a dark green to greenish brown homogenous liquid having a distinct odor of Lagundi, with grapes flavor and characteristic Lagundi taste.

WHAT IS IN THE MEDICINE?

Negacof syrup contains Lagundi (*Vitex negundo L. Fam Verbenaceae*) dried leaves. It is the only extensively researched and clinically proven natural medicine for cough and asthma in the Philippines. Lagundi (*Vitex negundo L.*) leaves is the main component of the Negacof. It is a large native shrub that grows in the Philippines.

WHAT IS THE MEDICINE USED FOR?

This product is used for the relief of mild to moderate cough due to common colds, flu and mild to moderate acute bronchitis. For relief of mild to moderate reversible bronchospasm in adults and children over 2 years with obstructive airway disease such as asthma and chronic bronchitis.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THE MEDICINE?

Adults: 10 mL (2 teaspoonful) 3 to 4 times daily. Dose may be increased as needed.
Children: 15 mg/kg/dose (or 0.25 mL/kg/dose) administered 3 times daily.
2 – 4 years: ½ teaspoon to 1 teaspoon 3 times daily. (10 to 15.5 kgs.)
4 – 6 years: 1 teaspoon 3 times daily. (15.5 to 20 kgs.)
6 – 12 years: 1 ½ teaspoonful – 2 teaspoonful 3 times daily.
Children over 40 kgs body weight: 2 teaspoonful 3 times daily.
Or as prescribed by the physician.

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

If you are allergic to any ingredient of the product. Safety among pregnant and lactating women has not been studied.

UNDESIRABLE EFFECTS

There are undesirable effects on the tablet.

WHAT OTHER MEDICINE OR FOODS SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

It may be taken with or without food. Take before or after meal.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, just give the next dose and the subsequent doses at the usual recommended schedule. Do not double the dose unless recommended by a doctor.

HOW SHOULD YOU KEEP THE MEDICINE?

- Keep the product out of reach and sight of children.
- Store at temperatures not exceeding 30°C
- Protect from light
- Always keep container tightly closed.

SIGNS AND SYMPTOMS OF OVERDOSE

There is no known signs and symptoms of overdose.

WHAT TO DO WHEN YOU HAVE USED MORE THAN THE RECOMMENDED DOSAGE?

In case of an overdose, consult a doctor or pharmacist immediately.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- Do not use after the expiry date on the label.
- Do not give more than the recommended dose.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

If any undesirable effect occurs and if symptoms persist.

AVAILABILITY:

Boston Round Amber Glass Bottle x 60 mL and 120 mL (Box of 1's)

DATE OF OF PATIENT INFORMATION LEAFLET (PIL): August 2018

FDA Registration no. : HMR-113

ADR Reporting Statement

"For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph"
Patient must seek medical attention immediately at the first sign of any adverse drug reaction

Manufactured by:

LA CROESUS PHARMA INC.
Brgy. Sto. Tomas Biñan, Laguna

Manufactured for:

MEDILYN DRUG
4 Dominga St. Mendez Rd., Baesa Quezon City

Under License from: University of the Philippines Manila (UPM)
National Integrated Research Program on Medicinal Plants (NIRPROMP)